



Review

# Diversity and Control of Spoilage Fungi in Dairy Products: An Update

Lucille Garnier <sup>1,2</sup> , Florence Valence <sup>2</sup> and Jérôme Mounier <sup>1,\*</sup>

<sup>1</sup> Laboratoire Universitaire de Biodiversité et Ecologie Microbienne (LUBEM EA3882), Université de Brest, Technopole Brest-Iroise, 29280 Plouzané, France; lucille.garnier@univ-brest.fr

<sup>2</sup> Science et Technologie du Lait et de l'Œuf (STLO), AgroCampus Ouest, INRA, 35000 Rennes, France; florence.valence-bertel@inra.fr

\* Correspondence: jerome.mounier@univ-brest.fr; Tel.: +33-(0)2-90-91-51-00; Fax: +33-(0)2-90-91-51-01

Received: 10 July 2017; Accepted: 25 July 2017; Published: 28 July 2017

**Abstract:** Fungi are common contaminants of dairy products, which provide a favorable niche for their growth. They are responsible for visible or non-visible defects, such as off-odor and -flavor, and lead to significant food waste and losses as well as important economic losses. Control of fungal spoilage is a major concern for industrials and scientists that are looking for efficient solutions to prevent and/or limit fungal spoilage in dairy products. Several traditional methods also called traditional hurdle technologies are implemented and combined to prevent and control such contaminations. Prevention methods include good manufacturing and hygiene practices, air filtration, and decontamination systems, while control methods include inactivation treatments, temperature control, and modified atmosphere packaging. However, despite technology advances in existing preservation methods, fungal spoilage is still an issue for dairy manufacturers and in recent years, new (bio) preservation technologies are being developed such as the use of bioprotective cultures. This review summarizes our current knowledge on the diversity of spoilage fungi in dairy products and the traditional and (potentially) new hurdle technologies to control their occurrence in dairy foods.

**Keywords:** spoilage; fungi; dairy products; control; diversity

## 1. Introduction

Since people began producing and storing food products, spoilage and food losses and waste became important issues for human with regards to food safety and security. Nowadays, up to one third of all food is spoiled or squandered before consumption, which represents about 1.3 billion tons per year [1]. These losses are the results of one or more problems occurring in the supply chain, from initial agricultural production down to the consumer level [1]. Concerning food spoilage, a food product can be physically, chemically, or microbiologically spoiled. Parasites, bacteria, and/or fungi are the main agents causing microbial spoilage. Certain parasites and bacteria are of public health concern because they are common contaminants of foods and are often responsible for food outbreaks. Nonetheless, being ubiquitous in nature, fungi are also very common in food. For a long time, besides altering food properties, they were not regarded as particularly detrimental to human health, although *Claviceps purpurea* has been related to ergotism outbreaks in the Middle Ages more than 200 years ago [2]. It is only in recent times that several mycotoxins produced by certain fungal species have been regarded as a major threat to human and animal health, especially in developing countries, being responsible for different adverse health effects [3].

Yeasts and molds are able to grow in a large variety of food including raw materials such as cereals, vegetables, fruits, meat, and milk, as well as processed products [4]. Given this large variety,

fungal spoilage leads to severe economic losses for food manufacturers. Indeed, it is estimated that between 5% and 10% of the world's food production is lost due to fungal deterioration [2]. Dairy products are less susceptible to fungal spoilage than other products, such as fruits or vegetables, because they are usually kept at refrigerated temperatures, are often made with heat-treated milk, and some of them are fermented products meaning that they may possess a competitive microbiota, have an acidic pH, and naturally contain organic acids. Nonetheless, a significant number of fungal species are able to survive and grow in dairy products. This astonishing adaptation capacity may be explained by the ability of fungi to utilize numerous substrates including carbohydrates, organic acids, proteins, and lipids that are present in milk and its derived products [5]. Moreover, these fungi are acidotolerant, xerotolerant, and/or psychrotolerant, and to some extent can tolerate chemical preservatives, which are sometimes added to increase product shelf-life. It is worth mentioning that several fungal species such as *Debaryomyces hansenii*, *Candida catenulata*, *Galactomyces geotrichum*, *Kluyveromyces marxianus*, *Mucor lanceolatus*, *Penicillium roqueforti*, *Penicillium camemberti*, or *Saccharomyces cerevisiae* are deliberately added as technological adjunct cultures to manufacture dairy products including kefir-type products and many kinds of cheese varieties [6–8]. Concerning undesirable species, their presence in dairy products may result in several types of food spoilage, e.g., visible growth of the fungus at the product surface, and the production of metabolites causing off-odors and flavors, as well as visible changes in color and/or texture [9]. In addition to organoleptic properties' deterioration, spoilage molds such as *Penicillium* and *Aspergillus* spp. can also produce mycotoxins [4,5,10,11]. In milk and dairy products, aflatoxin M1 (AFM1), which is produced by certain *Aspergillus* species, is the only mycotoxin for which maximum levels (0.05 ppb in the European Union (EU)) have been established. The occurrence of AFM1 in milk results from the conversion by dairy animals, fed on aflatoxin B1 (AFB1)-contaminated feedstuffs, of AFB1 to AFM1, which then pass to their urine and milk. Other mycotoxins such as ochratoxin A, citrinin, roquefortin C, mycophenolic, and cyclopiazonic acids have also been detected in cheeses at various concentrations [10]. Except for AFM1, intake of mycotoxins from dairy products is generally considered of limited importance compared to cereals and their derived products, and no human case of food poisoning related to mycotoxins has been documented so far.

Control of fungal spoilage is a major concern for industrials and scientists that are looking for efficient solutions to prevent and/or limit fungal growth or development in dairy products. Different traditional methods, also called traditional hurdle technologies, are implemented to control such contaminations including air treatment, cleaning and disinfection procedures, heat treatment, water activity reduction by brining, refrigeration, modified atmosphere packaging [12], as well as the use of chemical preservatives which are considered as food additives. The spoilage frequency and rate of many dairy foods can be reduced or retarded by the application of one or more of these treatments. However, fungal spoilage is still an issue for dairy manufacturers. Indeed, increasing fungal resistance toward heat treatments or chemical preservatives [13,14] and consumers' demand for more "natural" products, as well as legislation evolution have led industrial dairy producers to find complementary control approaches. This situation has led to the development of new (bio) preservation technologies such as the use of bioprotective cultures [15].

This review summarizes our current knowledge on spoilage fungi in dairy products with a special emphasis on their diversity, as well as the traditional and (potentially) new hurdle technologies to control their occurrence in dairy products.

## 2. Diversity of Spoilage Fungi in Dairy Products

With the use of multilocus sequencing of DNA regions with taxonomical interest, fungal taxonomy has undergone important changes during the last fifteen years. Many species have been reclassified and new phylogenetic species are also being regularly recognized within so-called "species complexes," in which members only harbor few differences in their morphological characters or are even morphologically undistinguishable from each other. For fungal identification, the internal transcribed spacer region (ITS) has been chosen as the best universal barcode [16]. Concerning yeast

identification, the D1/D2 domain of the 28 S rRNA gene is also widely used. It is worth mentioning that these genes are not always sufficient to identify isolates at the species level, and sequencing of other genes can be required. This is especially true for members of the *Aspergillus*, *Cladosporium*, *Penicillium*, and *Phoma* genera.

Over the years, the microbial diversity of milk and fermented dairy products has received considerable attention, and the development of next-generation sequencing (NGS) technologies [17–22] offers novel and rapid methods to characterize food ecosystems. However, except for raw milk, these techniques are not widely used to investigate fungal spoilage of dairy products because spoilage is generally the consequence of the outgrowth of a few species at the same time and because contamination can also occur on a distinct spot of the product surface. In addition, because read length is still limited with the currently used NGS technologies and because the taxonomic resolution of a single barcode marker can vary among taxa, it can be very tricky to identify operational taxonomic units (OTUs) at the species level. Nevertheless, these techniques offer new possibilities to investigate the potential sources of fungi in the dairy environment, which constitutes an important source of spoilage microorganisms.

### 2.1. Sources of Fungal Contamination

Fungal contamination of dairy foods can occur at different stages, from dairy farms to dairy processing units and at consumers' homes. Independent of the animal species, raw milk generally contains between 3 to 5 log 10 CFU·mL<sup>−1</sup> fungi with higher number of yeast cells than fungal spores [23,24]. As shown by a recent study [25], the stable and milking parlor environments at the farm are important sources of fungi in the milk. In addition, an important yeast source is the teat surface [25]. It should be mentioned that yeast growth during milk storage is rare, as yeasts are rapidly outnumbered by psychrotrophic bacteria such as *Pseudomonas* spp. [26,27]. Moreover, except for a few fungal species, yeast and molds are not heat-resistant and should be killed after pasteurization. Therefore, during manufacturing, fungal contamination generally occurs after milk heat-treatment. Mold spoilage is often due to airborne fungi because fungal spores are easily dispersed into the dairy plant air [28]. In a recent study undertaken in a Greek dairy plant, fungal counts of 362.3 CFU/m<sup>3</sup> consisting mainly of mold spores were reported in outdoor air samples while 69.8 CFU/m<sup>3</sup> and 266.2 CFU/m<sup>3</sup> were found in samples from two indoor locations [29]. Recent studies [28–33] investigated the fungal diversity of the dairy environment in different dairy plants, and mold contaminations were also shown to originate from the air. In the case of sliced cheese, the cheese rind itself can also be a source of spoilage fungi which are transferred during cheese slicing to the cheese slice surface and will grow during storage at the retailer or at the consumer's home [30]. In contrast, yeast causing spoilage generally originates from brine, surface, equipment, or ingredient contaminations [32,34–36], but can also be detected in the air [29]. The brine used for cheese salting/curing is one of the most significant sources of fungi. In one study [32], counts of 1.10<sup>9</sup> CFU/cm<sup>2</sup> were reported in a brining tank. Moreover, ingredients such as fruit preparations can also be important contamination sources for both yeast and molds [4,37]. In addition, certain fruit preparations such as blueberries and strawberries are even more at risk as they do not support extensive heat-treatment [38]. It is also worth mentioning that 50–100% fruit preparations including lemon cells, strawberries, and blueberries were recently reported to contain heat-resistant ascospore-forming fungi [39]. Finally, packaging materials may also be a source of spoilage molds, despite the fact that it has not been extensively studied.

Although there has been limited research on the subject, fungal contaminations are also likely to occur at the consumer's home after product opening. In a recent study on 586 surface samples collected from ~293 refrigerators in Italy [40], 15% and 5% of total samples contained fungal populations >1 log<sub>10</sub> CFU·cm<sup>−2</sup> and >2.5 log<sub>10</sub> CFU·cm<sup>−2</sup>, respectively. Besides refrigerator surfaces, cross-contamination from one product to another can also be responsible for fungal contamination. For example, mold- and smear-ripened cheeses as well as vegetables and fruits are potential yeast and mold contamination sources. Finally, the indoor air at a consumer's home is also a potent source of spoilage molds.

## 2.2. Spoilage Yeasts

Dairy food spoilage caused by yeast can result in visible alterations mainly due to their outgrowth on the product surface, such as the “toad skin” defect caused by *G. geotrichum* or browning reaction caused by *Yarrowia lipolytica*. The latter defect is due to the extracellular accumulation of homogentisic acid, an intermediate of tyrosine catabolism, capable of auto-oxidization and polymerization, leading to the formation of pyomelanin, a brown pigment [41,42]. Yeast spoilage may also lead to detectable but non-visible alterations resulting in off-odor and -flavor or texture alterations through the production of ethanol, CO<sub>2</sub>, and volatile organic compounds (primary and secondary alcohols, aldehydes, esters) as well as the production of lipolytic and proteolytic enzymes (glycolysis). One should keep in mind that the extent of spoilage depends on two parameters: the minimal spoilage level and the chemical spoilage index, which represent the concentrations of specific spoilage organism(s) and that of spoilage metabolites determined at the time of sensory rejection, respectively [43]. In relation to yeast spoilage in dairy products, little data are available on these values. It was defined that yeasty and fermented off-flavors were detected when yeasts grew at populations equal or above 10<sup>5</sup>–10<sup>6</sup> CFU/g [9].

It is also worth mentioning that certain yeasts species are able to produce biogenic amines (BA) including histamine and tyramine [44–46]. For example, *Y. lipolytica* was responsible for histidine, lysine, ornithine, phenylalanine, and tyrosine decarboxylation, leading to the production of putrescine, 2-phenylethylamine, tyramine, and cadaverine, respectively, in a traditional Italian cheese [44]. However, the main BA producers in dairy products are bacteria including lactic acid bacteria (LAB) and enterobacteria, particularly with the formation of histamine and tyramine [47]. Interestingly, it has already been shown that *G. geotrichum* was able to slightly degrade BA such as tyramine [48].

A significant number of studies have been performed to assess spoilage yeast diversity (Table 1). These contaminations can be due to one or several species. For example, a recent study showed that up to 14 different yeast species could be encountered in as low as six fresh cheese samples, including cases of multi-contaminations [49]. Until now, more than 60 yeast species have been identified as spoilage agents of dairy products (Table 1). They belong to the Ascomycota and Basidiomycota phylum represented by 20 and 10 genera, respectively. Among Ascomycota, all identified species belong to the Saccharomycotina subdivision and Saccharomycetes class, which is the only class of this subdivision. Among this class, the genus *Candida* is the most frequently reported, representing half of the Ascomycota diversity, and is also characterized by a high diversity at the species level. Indeed, 24 different *Candida* species have already been reported as responsible for dairy product spoilage. *Candida parapsilosis* is the most frequently isolated species, followed by *Candida lusitanae*, *Candida inconspicua*, and *Candida intermedia*. All these species can thus be considered common contaminants of dairy products. Table 1 also shows that dairy products spoiled by *Candida* are principally unripened products and hard or semi-hard cheeses such as Scamorza or Caciotta.

Besides *Candida* spp., *Debaryomyces*, *Kluyveromyces*, *Yarrowia*, *Galactomyces*, and *Saccharomyces* spp. are also frequent spoilers of fresh dairy products (fresh cheese, cream, and yoghurt) and heat-treated products (Table 1). The presence of spoilage yeasts in the latter products is principally due to post-heat-treatment contaminations, but certain yeast species show high resistance to pasteurization [50] (see Section 3.2.1). Contrary to the genus *Candida*, one or two species from each genus are considered dairy spoilage agents. Indeed, the *Kluyveromyces* genus is represented by *Kluyveromyces marxianus* and *Kluyveromyces lactis*. *Debaryomyces*, *Galactomyces*, *Saccharomyces*, and *Yarrowia* genera are represented by *D. hansenii*, *G. geotrichum*, *S. cerevisiae*, and *Y. lipolytica*, respectively. Other Ascomycota yeasts responsible for spoilage include *Meyerozyma* (formerly *Pichia*) and *Pichia*, *Geotrichum*, *Dekkera*, *Torulaspora*, *Wickerhamomyces*, *Blastodendron*, *Cyberlindnera*, *Kazachstania*, *Peterozyma*, *Priceomyces*, and *Torulopsis* spp., principally isolated from soft and unripened dairy products such as fresh cheese or yoghurt.

Among Basidiomycota, six genera and four genera belonging to the Agaromycotina and Pucciniomycotina subdivisions are regularly encountered in spoiled dairy products. Among Pucciniomycotina, the genus *Rhodotorula* (Microbotryomycetes class), represented by three different

species, i.e., *Rhodotorula diffluens*, *Rhodotorula glutinis*, and *Rhodotorula mucilaginosa*, is the most frequently cited and is essentially isolated from unripened products (Table 1). The latter genus is followed by *Cryptococcus* (Tremellomycetes class), represented by four species, i.e., *Cryptococcus humicola*, *Cryptococcus laurentii*, *Cryptococcus pseudolongus*, and *Cryptococcus minuta* isolated from unripened products (fresh cheese and butter/margarine) and soft cheese (Table 1). Other *Basidiomycota* genera include *Saitozyma*, *Naganishia*, *Cutaneotrichosporon*, *Sporobolomyces*, *Sporodiobolus*, *Trichosporon*, and *Vanjira* spp.

**Table 1.** Diversity of spoilage yeasts in dairy products.

Phylum	Species	Product Types	References
Ascomycota	<i>Atelosaccharomyces pseudotropicalis</i>	Yoghurt	[26]
	<i>Blastodendron arztii</i>	Fresh unripened cheese	[51]
	<i>Candida acidothermophilum</i>	Yoghurt	[26,36,52]
	<i>Candida apis</i>	Hard or semi-hard cheese	[53]
	<i>Candida beerwijckiae</i>	Hard or semi-hard cheese	[51]
		Yoghurt	[26]
	<i>Candida blankii</i>	Heat-treated milk and dairy product	[53]
	<i>Candida glabrata</i>	Soft and semi-soft cheese	[51]
	<i>Candida glabrosa</i>	Hard or semi-hard cheese	[54]
	<i>Candida inconspicua</i>	Buffalo, goat, or sheep cheese	[51]
		Fresh unripened cheese	[49,51]
		Soft and semi-soft cheese	[51]
	<i>Candida intermedia</i>	Hard or semi-hard cheese	[2,54]
		Other dairy products	[49]
		Yoghurt	[26]
	<i>Candida krusei</i>	Soft and semi-soft cheese	[26]
	<i>Candida lusitanae</i>	Buffalo, goat, or sheep cheese	[51]
		Fresh unripened cheese	[51]
		Soft and semi-soft cheese	[51]
		Yoghurt	[26,52]
	<i>Candida multigemmis</i>	Hard or semi-hard cheese	[2,54]
	<i>Candida mycoderma</i>	Buffalo, goat, or sheep cheese	[51]
	<i>Candida norvegica</i>	Soft and semi-soft cheese	[11]
	<i>Candida parapsilosis</i>	Fresh unripened cheese	[51]
		Hard or semi-hard cheese	[2,51,54]
		Soft and semi-soft cheese	[11,49,51]
		Yoghurt	[2,49]
	<i>Candida pseudoglabrosa</i>	Raw milk	[27]
	<i>Candida rugosa</i>	Soft and semi-soft cheese	[11]
	<i>Candida saitoana</i>	Yoghurt	[26]
	<i>Candida sake</i>	Fresh unripened cheese	[55]
		Hard or semi-hard cheese	[54]
	<i>Candida sphaerica</i>	Fresh unripened cheese	[26]
	<i>Candida stellata</i>	Hard or semi-hard cheese	[53]
		Yoghurt	[26,53]
	<i>Candida versatilis</i>	Yoghurt	[26,36]
	<i>Candida zeylanoides</i>	Fresh unripened cheese	[51]
		Hard or semi-hard cheese	[51]
		Soft and semi-soft cheese	[51]
	<i>Candida</i> sp.	Mold ripened cheese	[26]
	<i>Cyberlindnera jadinii</i>	Soft and semi-soft cheese	[11]
	<i>Debaryomyces hansenii</i>	Blue-veined cheese	[26]
		Buffalo, goat, or sheep cheese	[51]
		Fresh unripened cheese	[51,53,55]
		Hard or semi-hard cheese	[26,49,51]
		Heat-treated milk and dairy product	[26,52,53]
		Other dairy products	[52,53]
		Soft and semi-soft cheese	[11,51]
		Mold ripened cheese	[26]
		Yoghurt	[2,26,36,52,53]
	<i>Dekkera anomala</i>	Fresh unripened cheese	[55]



Table 1. Cont.

Phylum	Species	Product Types	References
	<i>Exophiala</i> sp. <i>Galactomyces geotrichum</i>	Yoghurt	[26]
		Fresh unripened cheese	[49]
		Butter and margarine	[2]
		Fresh unripened cheese	[2,49,55]
	<i>Geotrichum capitatum</i> <i>Geotrichum klebahnii</i> <i>Geotrichum</i> sp.	Heat-treated milk and dairy product	[2]
		Other dairy products	[49]
		Soft and semi-soft cheese	[51]
		Hard or semi-hard cheese	[51]
	<i>Kazachstania unispora</i> <i>Kluyveromyces lactis</i> <i>Kluyveromyces marxianus</i>	Buffalo, goat, or sheep cheese	[56]
		Raw milk	[57]
		Soft and semi-soft cheese	[11]
		Fresh unripened cheese	[49,55]
	<i>Meyerozyma guilliermondii</i>	Blue-veined cheese	[26]
		Fresh unripened cheese	[26,53]
		Hard or semi-hard cheese	[26,53,54]
		Heat-treated milk and dairy product	[26,53]
		Mold ripened cheese	[26]
		Other dairy products	[52]
		Raw milk	[27]
		Soft and semi-soft cheese	[49]
		Yoghurt	[2,26,36,53]
		Buffalo, goat, or sheep cheese	[51]
		Fresh unripened cheese	[49]
		Soft and semi-soft cheese	[11,51]
	<i>Naumovia dairenensis</i> <i>Peterozyma toletana</i> <i>Pichia fermentans</i> <i>Pichia membranifaciens</i>	Yoghurt	[49]
		Soft and semi-soft cheese	[11]
		Yoghurt	[36]
		Soft and semi-soft cheese	[11,49]
	<i>Pichia norvegensis</i> <i>Priceomyces carsonii</i> <i>Saccharomyces cerevisiae</i>	Fresh unripened cheese	[26]
		Soft and semi-soft cheese	[11]
		Soft and semi-soft cheese	[11]
		Hard or semi-hard cheese	[54]
		Blue-veined cheese	[26]
		Buffalo, goat, or sheep cheese	[51]
		Fresh unripened cheese	[51,53]
		Hard or semi-hard cheese	[53]
		Heat-treated milk and dairy product	[26]
		Other dairy products	[52]
		Soft and semi-soft cheese	[11,51]
		Yoghurt	[26,36,53]
	<i>Torulaspora delbrueckii</i> <i>Torulopsis</i> sp. <i>Wickerhomomyces anomalus</i>	Fresh unripened cheese	[51]
		Soft and semi-soft cheese	[11]
		Yoghurt	[26]
		Soft and semi-soft cheese	[11]
	<i>Yarrowia lipolytica</i>	Yoghurt	[2]
		Blue-veined cheese	[26]
		Buffalo, goat, or sheep cheese	[51]
		Butter and margarine	[53]
		Fresh unripened cheese	[26,55]
		Hard or semi-hard cheese	[2,49,51,54]
		Mold ripened cheese	[2,26]
		Soft and semi-soft cheese	[11,49]
		Yoghurt	[2]
		Butter and margarine	[2]
Basidiomycota	<i>Cryptococcus humicola</i>	Hard or semi-hard cheese	[2]
		Butter and margarine	[53]
	<i>Cryptococcus laurentii</i>	Fresh unripened cheese	[26,53]
		Fresh unripened cheese	[49]
	<i>Cryptococcus pseudolongus</i> <i>Cryptococcus</i> sp.	Heat-treated milk and dairy products	[52]
		Soft and semi-soft cheese	[11]
	<i>Cystobasidium minuta</i> <i>Cutaneotrichosporon curvatus</i> <i>Cutaneotrichosporon cutaneum</i>	Soft and semi-soft cheese	[11]
		Raw milk	[26]
		Hard or semi-hard cheese	[51]
		Fresh unripened cheese	[51]

Table 1. Cont.

Phylum	Species	Product Types	References
	<i>Naganishia albida</i>	Butter and margarine	[53]
		Fresh unripened cheese	[53]
		Other dairy products	[53]
		Yoghurt	[53]
	<i>Rhodotorula diffluens</i>	Butter and margarine	[53]
		Fresh unripened cheese	[53]
		Hard or semi-hard cheese	[26,53]
		Heat-treated milk and dairy products	[26,52,53]
		Yoghurt	[2,26,53]
	<i>Rhodotorula glutinis</i>	Butter and margarine	[53]
		Fresh unripened cheese	[53]
		Other dairy products	[52,53]
		Soft and semi-soft cheese	[11]
	<i>Rhodotorula mucilaginosa</i>	Butter and margarine	[53]
		Fresh unripened cheese	[51,53]
		Other dairy products	[52,53]
		Soft and semi-soft cheese	[49]
		Yoghurt	[2,36]
	<i>Rhodotorula sp.</i>	Heat-treated milk and dairy products	[26]
	<i>Saitozyma flava</i>	Fresh unripened cheese	[53]
		Hard or semi-hard cheese	[53]
		Heat-treated milk and dairy products	[26,53]
		Other dairy products	[53]
	<i>Sporobolomyces roseus</i>	Fresh unripened cheese	[26]
	<i>Sporodiobolus salmonicolor</i>	Hard or semi-hard cheese	[49]
	<i>Trichosporon asahii</i>	Soft and semi-soft cheese	[49]
	<i>Vanrija humicola</i>	Fresh unripened cheese	[51]

### 2.3. Molds in Dairy Products

Dairy product spoilage by molds also consists in visible alterations due to the outgrowth of a thallus on the product surface. In vacuum-packed cheeses such as cheddar cheese, another defect called thread mold defect is sometimes encountered and is characterized by mold growth in folds and wrinkles of the plastic film in which the cheese is packaged [54]. Mold spoilage can also lead to the formation of off-flavors. For example, *Moniliella suaveolens* and *Cladosporium herbarum* were shown to produce ketones causing rancidity defect in butter while *Penicillium glabrum*, *Penicillium expansum*, *Penicillium chrysogenum*, and *Cladosporium cladosporioides* produced off-flavors including 2-methylisoborneol and geosmin which possess strong musty and earthy notes [2]. Interestingly, a study [58] showed that certain fungal species were able to metabolize sorbate salts (used as preservatives) to produce metabolites such as *trans*-1,3-pentadiene or *trans*-piperylene, resulting in the formation of plastic-like or “kerosene” off-flavors.

Molds responsible for dairy product spoilage are highly diverse at the genus and species levels (Table 2). Indeed, up to 100 mold species have been identified so far as being responsible for dairy product spoilage. Most species belong to the Ascomycota and Mucoromycota phyla (Table 2). Both Ascomycota and Mucoromycota have been principally isolated from hard or semi-hard cheeses.

Among Ascomycota, all identified spoilage molds (88 species distributed in 26 genera) belong to the Pezizomycotina subdivision, with the exception of *Lecanicillium lecanii* [59]. Within the Pezizomycotina subdivision, the Eurotiomycetes class is the most represented (nine genera and 57 species), followed by the Dothideomycetes (10 genera and 20 species), the Sordariomycetes (11 species of six genera) and the Leotiomycetes (one species). Among Eurotiomycetes, *Penicillium* is by far the most frequently reported spoilage genus with ~40 species, followed by *Aspergillus* (10 species). Other genera include *Byssoschlamys*, *Eupenicillium*, *Eurotium*, *Exophiala*, *Hamigera*, *Neosartorya*, *Rhinochlaidiella*, and *Talaromyces* spp.

*Penicillium* spp. are mainly isolated from hard and semi-hard cheeses but are also found in all other product types including fresh, blue-veined, mold-ripened, soft- and semi-soft cheeses, butter, yoghurt, milk as well as in buffalo, goat, or ewe's milk cheeses. Based on the classification system recently proposed by Houbraeken and Samson [60], *Penicillium* spp. responsible for spoilage mainly belong to the Fasciculata section (11 species) but also to the Chrysogena (three species), Biverticillium (four species), Brevicompacta (three species) and Aspergilloides (two species) sections. Among the Fasciculata section, *Penicillium commune*, *Penicillium solitum*, *Penicillium crustosum*, and *Penicillium verrucosum* are frequently reported, while within the Chrysogena section, *P. chrysogenum*, *Penicillium nalgiovense*, and *Penicillium griseofulvum* are quite common. Within Brevicompacta, *Penicillium brevicompactum* is frequently reported, in addition to *Penicillium bialowiezense* [49] and *Penicillium spathulatum* [61]. Nonetheless, Samson et al. [62] underlined that *P. bialowiezense* could have been misidentified in previous studies with the closely related species *P. brevicompactum*. The *Penicillium* species mentioned above are adaptive to low temperatures, and several of them are considered xerophilic species [2], therefore, their occurrence in spoiled dairy products is not surprising. *Penicillium* spp. is also well-adapted to the cheese environment and has been associated with cheese manufacturing for centuries. In addition, a recent study highlighted the presence of not previously mentioned xerophilic species such as *Penicillium charlesii*, *Penicillium fellutanum*, *Penicillium adametzioides*, and *Penicillium antarcticum* [49].

Fungi from the Dothideomycetes class are principally isolated from hard and semi-hard cheeses and constitute the second most frequently identified class after Eurotiomycetes. Among this class, *Cladosporium*, with seven species distributed in three different *Cladosporium* complexes, is the most frequently reported. *Cladosporium* spp. grow slowly but are very common airborne fungi and are quite psychrotolerant and xerotolerant [2].

Among the Mucoromycotina sub-phylum, three different genera have already been isolated from contaminated dairy products, i.e., *Mucor* (four species), *Rhizopus* (one species), and *Thamnidium* (one species), all belonging to the Mucorales order. As for other genera belonging to the Ascomycota phylum, these genera were mostly isolated from spoiled cheeses [56,62,63].

Molds can also contaminate heat-treated dairy products. Indeed, species belonging to *Aspergillus*, *Byssochlamys*, *Cladosporium*, *Eupenicillium*, *Fusarium*, *Hamigera*, *Neosartorya*, *Penicillium*, and *Talaromyces* genera have been isolated from pasteurized milk, cream cheese, and heat-treated dairy beverages [2,49,57]. The presence of mold in heat-treated milk or dairy products might be due to post-processing contaminations during bottling or packaging [57] or to the heat-resistance of mold spores. Indeed, Pitt and Hocking (2009) isolated heat-resistant species such as *Byssochlamys nivea* (anamorph *Paecilomyces niveus*), *Eupenicillium brefeldianum*, *Hamigera avellanea*, and *Neosartorya fischeri* (anamorph *Aspergillus fischeri*) from cream cheese, as well as *B. nivea*, *N. fischeri*, and *Talaromyces macrosporus* from heat-treated dairy beverage [2]. In a recent study, it was shown that the dormancy of *Talaromyces* and *Neosartorya* spp. ascospores was broken during thermal processing, for which heat-treatment was not optimized for ascospore inactivation [64].

In addition, fungal spoilage of dairy products obtained by Ultra-High Temperature (UHT) treatment can also occur, but it is generally due to post-processing contaminations. For example, *Fusarium oxysporum* was isolated on several occasions from UHT-flavored milk beverages [2].



**Table 2.** Diversity of spoilage filamentous fungi isolated from contaminated dairy products.

Phylum	Genera	Species	Product Types	References
Ascomycota	<i>Acremonium</i> <i>Alternaria</i>	Nd *	Buffalo, goat, or sheep cheese	[56]
		<i>Alternaria alternata</i>	Blue-veined cheese	[63]
		Nd *	Hard or semi-hard cheese	[54,65]
			Hard or semi-hard cheese	[66]
	<i>Aspergillus</i> section <i>Aspergillus</i>	<i>Aspergillus glaucus</i>	Raw milk	[25]
			Heat-treated milk	[57]
	<i>Aspergillus</i> section <i>Candidi</i>	<i>Aspergillus candidus</i>	Raw milk	[57]
			Buffalo, goat, or sheep cheese	[56]
	<i>Aspergillus</i> section <i>Circumdati</i>	<i>Aspergillus ochraceus</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[67]
	<i>Aspergillus</i> section <i>Flavi</i>	<i>Aspergillus flavus</i>	Heat-treated milk	[57]
			Raw milk	[57]
	<i>Aspergillus</i> section <i>Fumigati</i>	<i>Aspergillus tamarii</i> <i>Aspergillus fumigatus</i>	Blue-veined cheese	[63]
			Blue-veined cheese	[63]
			Hard or semi-hard cheese	[65]
			Heat-treated milk	[57]
	<i>Aspergillus</i> section <i>Nidulantes</i>	<i>Aspergillus sydowii</i> <i>Aspergillus versicolor</i>	Raw milk	[57]
			Yoghurt	[2]
			Blue-veined cheese	[63]
			Hard or semi-hard cheese	[59,68]
	<i>Aspergillus</i> section <i>Nigri</i>	<i>Aspergillus niger</i>	Semi-soft cheese	[4,59]
			Buffalo, goat, or sheep cheese	[56]
	<i>Aspergillus</i>	Nd *	Blue-veined cheese	[63,67]
			Hard or semi-hard cheese	[54,66]
	<i>Aureobasidium</i>	<i>Aureobasidium pullulans</i>	Raw milk	[25]
			Hard or semi-hard cheese	[69]
	<i>Bipolaris</i>	<i>Bipolaris australiensis</i>	Blue-veined cheese	[63]
	<i>Botrytis</i>	<i>Botrytis cinerea</i>	Hard or semi-hard cheese	[54]
	<i>Byssoschlamys</i>	<i>Byssoschlamys nivea</i>	Heat-treated milk	[2]
	<i>Cladosporium</i> <i>cladosporioides</i> complex	<i>Cladosporium</i> <i>cladosporioides</i>	Heat-treated milk	[2]
			Blue-veined cheese	[63]
			Buffalo, goat, or sheep cheese	[56]
			Butter and margarine	[2]
	<i>Cladosporium</i> <i>herbarum</i> complex	<i>Cladosporium phyllophilum</i> <i>Cladosporium butyri</i>	Hard or semi-hard cheese	[2,54]
			Hard or semi-hard cheese	[54]
	<i>Cladosporium</i> <i>sphaerospermum</i> complex	<i>Cladosporium herbarum</i> <i>Cladosporium macrocarpum</i>	Butter and margarine	[49]
			Butter and margarine	[2]
	<i>Cladosporium</i>	<i>Cladosporium halotolerans</i> <i>Cladosporium sphaerospermum</i>	Heat-treated milk	[49]
			Hard or semi-hard cheese	[49,54]
	<i>Didymella</i>	<i>Didymella pinodella</i>	Hard or semi-hard cheese	[59,66]
			Semi-soft cheese	[59]
	<i>Epicoccum</i>	<i>Epicoccum nigrum</i>	Fresh unripened cheese	[49]
			Other dairy product	[49]
	<i>Eupenicillium</i>	<i>Eupenicillium brefeldianum</i>	Hard or semi-hard cheese	[54,69]
			Heat-treated milk	[2]
	<i>Eurotium</i>	<i>Eurotium herbariorum</i>	Heat-treated milk	[2]
			Hard or semi-hard cheese	[65]
	<i>Exophiala</i>	Nd *	Raw milk	[25,68]
			Cream cheese	[49]
	<i>Fusarium</i>	<i>Fusarium avenaceum</i> <i>Fusarium oxysporum</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[54,59]
			Semi-soft cheese	[59]
			Heat-treated milk	[2]
		<i>Fusarium solani</i>	Hard or semi-hard cheese	[59]
			Semi-soft cheese	[59]
		<i>Fusarium verticillioides</i>	Hard or semi-hard cheese	[67]
			Hard or semi-hard cheese	[66]
		Nd *	Heat-treated milk	[57]
			Raw milk	[57]
	<i>Gliocladium</i>	<i>Gliocladium roseum</i>	Hard or semi-hard cheese	[54]
	<i>Hamigera</i>	<i>Hamigera avellanea</i>	Heat-treated milk	[2]
	<i>Lecanicillium</i>	<i>Lecanicillium lecanii</i>	Hard or semi-hard cheese	[59]
			Semi-soft cheese	[59]
	<i>Naumovia</i>	<i>Naumovia dairenensis</i>	Soft cheese	[11]
	<i>Neosartorya</i>	<i>Neosartorya fischeri</i>	Heat-treated milk	[2]
	<i>Penicillium</i> section <i>Aspergilloides</i>	<i>Penicillium glabrum</i>	Butter and margarine	[2]
			Hard or semi-hard cheese	[2,54,70]
			Fresh unripened cheese	[2]
			Yoghurt	[2]

Table 2. Cont.

Phylum	Genera	Species	Product Types	References
		<i>Penicillium spinulosum</i>	Hard or semi-hard cheese	[65]
	<i>Penicillium</i> section <i>Biverticillium</i>	<i>Penicillium funiculosum</i>	Buffalo, goat, or sheep cheese	[56]
		<i>Penicillium minioluteum</i>	Hard or semi-hard cheese	[69]
		<i>Penicillium purpureogenum</i>	Hard or semi-hard cheese	[68]
		<i>Penicillium rugulosum</i>	Hard or semi-hard cheese	[54,59]
			Semi-soft cheese	[59]
	<i>Penicillium</i> section <i>Brevicompacta</i>	<i>Penicillium bialowiezense</i>	Hard or semi-hard cheese	[49]
			Yoghurt	[49]
		<i>Penicillium brevicompactum</i>	Hard or semi-hard cheese	[2,49,54,59,65,69,70]
			Semi-soft cheese	[59]
			Buffalo, goat, or sheep cheese	[56]
	<i>Penicillium</i> section <i>Sclerotiora</i>	<i>Penicillium adametzioides</i>	Fresh unripened cheese	[49]
	<i>Penicillium</i> section <i>Canescentia</i>	<i>Penicillium antarcticum</i>	Hard or semi-hard cheese	[49]
		<i>Penicillium arenicola</i>	Buffalo, goat, or sheep cheese	[56]
		<i>Penicillium canescens</i>	Hard or semi-hard cheese	[68]
		<i>Penicillium novae-zeelandiae</i>	Hard or semi-hard cheese	[54]
	<i>Penicillium</i> section <i>Charlesii</i>	<i>Penicillium charlesii</i>	Fresh unripened cheese	[49]
			Heat-treated milk	[49]
		<i>Penicillium dierckxii</i>	Fresh unripened cheese	[49]
	<i>Penicillium</i> section <i>Chrysogena</i>	<i>Penicillium chrysogenum</i>	Butter and margarine	[2]
			Blue-veined cheese	[63]
			Hard or semi-hard cheese	[54,59,62,65,69,70]
			Semi-soft cheese	[59]
			Buffalo, goat, or sheep cheese	[56]
			Fresh unripened cheese	[2]
			Yoghurt	[2]
		<i>Penicillium griseofulvum</i>	Hard or semi-hard cheese	[68]
		<i>Penicillium nalgiovense</i>	Hard or semi-hard cheese	[4,49,59]
			Semi-soft cheese	[4,59]
			Buffalo, goat, or sheep cheese	[56]
	<i>Penicillium</i> section <i>Citrina</i>	<i>Penicillium steckii</i>	Hard or semi-hard cheese	[69]
	<i>Penicillium</i> section <i>Exilicaulis</i>	<i>Penicillium corylophitum</i>	Hard or semi-hard cheese	[65]
	<i>Penicillium</i> section <i>Fasciculata</i>	<i>Penicillium aurantiogriseum</i>	Hard or semi-hard cheese	[65,69]
			Buffalo, goat, or sheep cheese	[56]
			Yoghurt	[2]
		<i>Penicillium camembertii</i>	Hard or semi-hard cheese	[68]
		<i>Penicillium commune</i>	Other dairy products	[49]
			Hard or semi-hard cheese	[2,4,49,54,59,65,69,70]
			Mold ripened cheese	[2]
			Semi-soft cheese	[4,59]
			Buffalo, goat, or sheep cheese	[56]
			Fresh unripened cheese	[2]
		<i>Penicillium crustosum</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[54,59,62,65,69,70]
			Semi-soft cheese	[59]
			Buffalo, goat, or sheep cheese	[56]
		<i>Penicillium discolor</i>	Hard or semi-hard cheese	[49,59]
			Semi-soft cheese	[59]
		<i>Penicillium echinulatum</i>	Hard or semi-hard cheese	[59,65,68,69]
			Semi-soft cheese	[59]
		<i>Penicillium hirsutum</i>	Buffalo, goat, or sheep cheese	[56]
		<i>Penicillium nordicum</i>	Hard or semi-hard cheese	[49]
		<i>Penicillium palitans</i>	Fresh unripened cheese	[49]
			Hard or semi-hard cheese	[49,65,69]
		<i>Penicillium solitum</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[2,49,54,59,65,69]
			Other dairy products	[49]
			Semi-soft cheese	[4,59]
			Yoghurt	[49]
		<i>Penicillium verrucosum</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[54,59,68,69]
			Mold ripened cheese	[2]
			Semi-soft cheese	[4,59]
		<i>Penicillium viridicatum</i>	Hard or semi-hard cheese	[2,54,65,67–69]
	<i>Penicillium</i> section <i>Paradoxa</i>	<i>Penicillium atramentosum</i>	Blue-veined cheese	[63]
			Hard or semi-hard cheese	[59]
			Semi-soft cheese	[59]
	<i>Penicillium</i> section <i>Penicillium</i>	<i>Penicillium expansum</i>	Butter and margarine	[2]
			Hard or semi-hard cheese	[2,65,68,69]
	<i>Penicillium</i> section <i>Lanata-Divaricata</i>	<i>Penicillium oxalicum</i>	Buffalo, goat, or sheep cheese	[56]
	<i>Penicillium</i> section <i>Roquefortum</i>	<i>Penicillium simplicissimum</i>	Hard or semi-hard cheese	[57]
		<i>Penicillium roquefortii</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[2,54,59,65,68,69]
			Mold ripened cheese	[2]
			Semi-soft cheese	[4,59]
	<i>Penicillium</i>	Nd *	Buffalo, goat, or sheep cheese	[56]

Table 2. Cont.


Phylum	Genera	Species	Product Types	References
Basidiomycota Zygomycota	<i>Phaeramularia</i> <i>Phoma</i>	Nd *	Hard or semi-hard cheese	[54,66]
			Heat-treated milk	[57]
			Raw milk	[25,57]
		Nd *	Hard or semi-hard cheese	[54]
		<i>Phoma glomerata</i>	Hard or semi-hard cheese	[2,70]
		<i>Phoma nebulosa</i>	Hard or semi-hard cheese	[69]
		Nd *	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[2,54,65,69]
		Nd *	Blue-veined cheese	[63]
		<i>Sarocladium strictum</i>	Hard or semi-hard cheese	[69]
	<i>Rhinocladia</i> <i>Sarocladium</i> <i>Scopulariopsis</i>	<i>Scopulariopsis brevicaulis</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[51,59]
			Semi-soft cheese	[4,59]
		Nd *	Hard or semi-hard cheese	[54,67]
		<i>Talaromyces</i>	Heat-treated milk	[2]
		<i>Trichoderma</i>	Hard or semi-hard cheese	[65]
		<i>Ulocladium</i>	Hard or semi-hard cheese	[69]
		<i>Wallenia sebi</i>	Raw milk	[25]
		<i>Mucor</i>	Hard or semi-hard cheese	[49,69]
			Yoghurt	[2]
		<i>Mucor hiemalis</i>	Buffalo, goat, or sheep cheese	[56]
			Hard or semi-hard cheese	[2,65,70]
			Yoghurt	[2]
		<i>Mucor plumbeus</i>	Hard or semi-hard cheese	[65,69]
		<i>Mucor racemosus</i>	Hard or semi-hard cheese	[49,59,65,68,69]
			Semi-soft cheese	[59]
			Yoghurt	[2]
		<i>Rhizopus</i>	Blue-veined cheese	[63]
			Buffalo, goat, or sheep cheese	[56]
	<i>Thamnidium</i>	<i>Thamnidium elegans</i>	Hard or semi-hard cheese	[49]

\* Nd: Not determined.

### 3. Prevention and Control of Fungal Spoilage in Dairy Products

Today, fungal spoilage continues to be a major challenge for industrials and, in modern practices, both preventive and control approaches are usually combined to reduce its incidence in dairy foods (Figure 1). Preventive methods can be defined as methods which can avoid contamination or recontamination during product processing, including packaging in aseptic conditions, use of air filtration system, and good manufacturing practices such as good hygienic practices and Hazard Analysis and Critical Control Points (HACCP) system implementation [71]. Control methods involve the use of methods which will either slow down or inhibit microbial growth, such as the addition of chemical preservatives, use of modified atmosphere packaging, and low temperature storage, as well as methods which will inactivate fungi such as heat and high-pressure treatments [71].

When it comes to choosing the most appropriate preventive and control methods, different factors must be taken into account. One major factor is the characteristics of the product to be preserved (composition,  $a_w$ ) and those of the microorganisms of concern as well as the sanitation conditions along the manufacturing process and product storage conditions. Another important factor is also consumer perception and acceptance of food preservation methods as well as their impact on hygiene and safety and their associated cost [72,73]. That is probably the reason why, among emerging methods to combat fungal spoilage, the use of bioprotective cultures and fermentates to decrease the use of chemical preservatives is gaining more and more attention [74].



**Figure 1.** Preventive and control methods used in dairy products for preventing, inactivating, retarding, or inhibiting growth of spoilage fungi.

### 3.1. Preventive Methods

#### 3.1.1. Good Manufacturing and Distribution Practices

The success of any food preservation method depends on low initial levels of contamination, which in turn are achieved through the application of appropriate cleaning, sanitation, decontamination, and hygienic practices [73]. Dairy product manufacturing and packaging must also be done in the most hygienic and aseptic conditions possible to reduce the risk of biological contamination including that of fungal contamination [75,76]. In a code of good manufacturing and distribution practices (GMDPs), two main points are essential: (1) selecting good quality raw materials and monitoring the manufacturing process to control microbes while preventing cross-contamination; and (2) avoiding or retarding microbial growth [77,78]. GMDPs are an indispensable part of every food quality system. This code relies on the use of good hygiene practices and on the implemented hazard analysis critical control point (HACCP) systems from the production to the consumption of dairy products. In accordance with the *Codex Alimentarius* indications and Regulation 852/2004/CE, HACCP systems identify, evaluate, and control hazards that threaten food hygiene, in particular taking into account the microbiological risks [79,80]. Thus, the microbiological quality of dairy products is controlled at each step of the production process, from the raw materials to the finished product [81]. Prerequisite programs, such as Total Quality Management (TQM), a method including planning, organizing, and understanding each activity and involving each individual at each level, or Just-In-Time (JIT), a program designed to reduce waste by reducing flow times within the production system as well as response times from suppliers and to customers, are useful to improve product

quality [81,82]. These programs have significant costs and require important efforts, but will produce return in investment in the long term.

### 3.1.2. Air Filtration and Decontamination Systems

With specific regard to the fungal contamination risk, potential contamination sources include ambient air. Therefore, efficient air-filtration systems should be in place to reduce spore counts into the air where the product is vulnerable. For example, Beletstiotis et al. [29] showed that the operation of High Efficiency Particulate Air (HEPA) filters class 10.000, designed to withhold fungal spores, allowed a decrease in fungal loads by 30 times in the indoor air of a dairy plant. The direction of air flows and location of outlets should also be carefully examined in sensitive zones. Moreover, the control of air pressurization can be applied to prevent air flow from the dirtiest to cleanest areas. Cleanroom technologies (aseptic or ultraclean technologies) including minienvironments that protect equipment such as filling and packaging equipment, transport system tunnels, or protective enclosures can also be used.

For air decontamination, gaseous ozone was successfully applied for disinfecting empty cheese ripening and storage facilities with up to a 10-fold reduction in mold and yeast counts depending on the dose and treatment time [83,84], see for a review Varga and Szigeti, 2016 [85]. Finally, molds and yeasts are able to grow in humid zones (walls, ceiling, and floor) and on the equipment if these are not properly cleaned and disinfected.

Various disinfectants such as alcohol, peracetic acid, iodophors, aldehydes, quaternary amine compounds, chlorine-based agents, or hydrogen peroxide have been used in the dairy industry against fungal contaminations. Nonetheless, fungal response to such cleaning agents varies with disinfectant type and concentration. For example, Korukluoglu et al. [86] showed that *Aspergillus niger* was sensitive to alcohol but not to peracetic acid; moreover, *A. niger* was resistant to 0.5% iodophor but sensitive at 1%. In addition, Bundgaard-Nielsen and Nielsen (1995) demonstrated the existence of intraspecific variation in tolerance to quaternary ammonium compounds, especially in *P. commune* and *Penicillium nalgiovense* [87]. Finally, Martin et al. [88] revealed that the combination of hydrogen peroxide and formic acid had a synergistic action against *Scopulariopsis brevicaulis* and *G. geotrichum*, thus showing that a combination of different disinfectants is a possible solution to prevent fungal contaminations. More work is required to investigate the efficiency of disinfectant agents against fungi in the dairy product context, while taking into account the continuously evolving legislation on their use.

## 3.2. Control Methods

### 3.2.1. Inactivation Treatments

Concerning heat-treatments, thermization, pasteurization, and UHT sterilization are intended to partially or completely reduce milk microbial load while altering as little as possible milk organoleptic and nutritional properties [89]. During such heat-treatment, the product is subjected to a high temperature for a short period of time. In the EU, thermization, pasteurization, and UHT correspond to treatments at 57–68 °C for a minimum of 15 s, 71.7 °C for 15 s, and at least 135 °C for a few seconds, respectively [90]. According to Sakkas et al. [91], the efficacy of heat treatments are related to the temperature–time combinations, heating method utilized, and milk pre-treatment conditions (filtration, homogenization, standardization, etc.). Concerning milk thermization, the main objective is to reduce the population of spoilage microorganisms and to inactivate enzymes while minimizing heat damage to milk components. Milk pasteurization is much more efficient than thermization, but not all microorganisms can be destroyed and this treatment has to be followed by a rapid chilling (4 °C) to slow down subsequent microbial growth. Finally, UHT sterilization, used for milk and dairy drinks, is itself obtained by applying a continuous heat-flow for a short time-period which allows for the total destruction of microorganisms followed by milk aseptic packaging [87].



As mentioned earlier, certain fungal species, because of their considerable heat-resistance, are able to spoil heat-treated dairy products, in particular pasteurized fruit-based products. Heat-resistant species include *Byssoschlamys spectabilis*, *N. fischeri*, *H. avellanea*, and *Talaromyces bacillisporus* [92,93]. This feature is due to their heat-resistant ascospores which are activated when heat-treatments are not high enough to inactivate them. As an example, *N. fischeri* and *T. bacillisporus* were shown to possess decimal reduction times at 85 °C ( $D_{85}$ ) ranging from 15.0 to 30.1 min, and from 11.9 and 15.5 min, respectively [39,94]. *N. fischeri* and *T. bacillisporus* showed  $D_{85}$  ranging from 47 to 75 min in *N*-(2-Acetamido)-2-aminoethanesulfonic acid buffer as well as  $D_{90}$  ranging from 7.5 to 12.7 min and  $D_{95}$  ranging from 0.56 to 0.67 min in blueberry juice [95]. They showed  $D_{90}$  from 2.0 to 7.6 min and  $D_{95}$  from 1.7 to 2.3 min in juices. *N. fischeri* and *T. bacillisporus* showed  $D_{87}$  from 11.1 to 66.7 min,  $D_{90}$  from 4.7 to 13.5 min, and  $D_{95}$  from 0.43 to 1.52 min and ranged between 44.4 and 60.9 min at 82 °C and 2.7 and 4.1 min at 88 °C. As another example, *H. avellanea* ascospores are activated but not destroyed after heat-treatment for 20 min at 87 °C, 5 min at 90 °C, and 1 min at 95 °C, showing the importance of time-temperature conditions treatment [93].

Other techniques such as high-pressure (HP) treatments of dairy products can also be used to inactivate yeast cells and mold spores (among other microorganisms) and extend product shelf-life [95]. Mainville et al. [96] and Reps et al. [97] showed that yeasts from kefir were completely inactivated at 400 MPa for 15 min or 30 min without significant changes in protein and lipid structure of the product. Moreover, Evert-Arriagada et al. [98] recently demonstrated that HP processing under industrial conditions could extend the shelf-life of a commercial starter-free fresh cheese. Pressurized cheeses were firmer and more yellow than control cheeses, but these changes did not affect the panel preference for pressurized cheese. It is worth mentioning that there is an increasing number of HP-treated dairy products available on the market as this technology meets consumer's expectations for more "natural" and less-heavily processed food products.

Pulsed electric fields (PEF) treatment consists of pulsed electric fields of strong intensity (15 to 50 kV/cm) for only few seconds [99]; this represents another emerging technology for microbial inactivation, including that of yeast and molds, with potential application in fluid dairy foods (see the recent reviews of Buckow et al. [100] and Wan et al. [101]).

### 3.2.2. Temperature Control

Temperature control is critical for dairy food quality and shelf-life. Cold temperatures are used to minimize microbial growth in raw milk until processing and to extend the shelf-life of non-sterile dairy foods. Low temperatures, or even frozen storage, however, do not eliminate microorganisms [73]. Milk and dairy products are generally stored between 0 °C and 4 °C and at temperatures ranging from 4 to 10 °C at the consumer's home. However, most fungi are psychrotrophic and are thus able to grow at such temperatures. In a recent study, Gougouli et al. [102] evaluated the effect of temperature and inoculum size on the growth of one isolate each from 12 fungal species during yoghurt storage. The ability to grow at refrigerated temperature was species-dependent with estimated minimal growth temperatures of −6.7 and 9.6 °C for *P. commune* and *A. niger*, respectively. In addition, Gougouli et al. [102] also developed and validated a probabilistic model to predict the appearance of visible mycelia originating from single spores which could be used for choosing adequate conditions for challenge tests. It should be noted that besides storage temperature and the spoilage organism itself, fungal spoilage susceptibility depends on the spoiler initial population and the time required to form a visible thallus on the product surface, which can be predicted using predictive mycology tools.

### 3.2.3. Modified Atmosphere Packaging

Modified atmosphere packaging (MAP) is a commonly used method to control fungal spoilage and thus to extend dairy food shelf-life. MAP consists of the replacement of air by a defined gas mixture. In MAP, active agents such as carbon dioxide [103] or oxygen scavengers [104] can also

be utilized. During storage, passive modifications of the headspace gases can occur, resulting from product respiration and gas transfer through the film.

As extensively reviewed by Nguyen Van Long et al. [105] on a large number of fungal species, lowering the oxygen level below 1% will result in the partial growth inhibition of strict aerobic fungi, while it will slow down facultative anaerobes' growth as it enables O<sub>2</sub> use as a final electron acceptor. However, for many species including *Penicillium* and *Mucor* spp., only 0% O<sub>2</sub> concentration can totally inhibit growth [105]. It is worth mentioning that such an O<sub>2</sub> level will favor the growth of facultative or strict anaerobes and that it can have a detrimental effect on products such as mold-ripened cheeses, for which a certain oxygen level is necessary to maintain their organoleptic properties [106]. An increase of the CO<sub>2</sub> level in package atmosphere can have an inhibitory effect, thus further extending product shelf-life [105]. In their extensive review, Nguyen Van Long et al. [105] indicated that, independent of the fungal species, a CO<sub>2</sub> level above 50% was necessary to achieve fungal growth inhibition and that, depending on the fungal species, 50–90% were necessary to completely inhibit fungal growth in solid matrices. Three mechanisms have been described to explain the CO<sub>2</sub> antimicrobial effect. The first mechanism is the displacement of some or all of the O<sub>2</sub> available for growth of aerobic microorganisms. The second one is a pH decrease resulting from CO<sub>2</sub> dissolution into the product and carbonic acid formation. Carbonic acid will then behave as a weak organic acid provoking disturbances in pH homeostasis [107,108]. Nonetheless, high CO<sub>2</sub> concentration may sometimes affect product organoleptic properties. For example, high CO<sub>2</sub> levels (100%) led to very negative effects in the sensory qualities of cottage cheese, whereas for others, such as cheddar, packaging with 100% CO<sub>2</sub> is commonly used in dairy industries [109,110].

MAPs are often used for cheeses and different gas compositions have been suggested [109,111,112]. For example, in commercial sliced cheddar cheese, *P. commune* did not grow in atmospheres composed either of 20%, 40%, and 60% CO<sub>2</sub> and less than 0.5% O<sub>2</sub> [113]. *Eurotium chevalieri* and *Xeromyces bisporus* could not grow under atmosphere containing 80% CO<sub>2</sub> and 20% O<sub>2</sub> during incubation for 60 days, whereas *Mucor plumbeus* development was observed after 15 days [111]. Facultative anaerobic fungi are also susceptible to high CO<sub>2</sub> levels [113–115]. As an example, a 60% CO<sub>2</sub>/40% N<sub>2</sub> modified atmosphere was effective for inhibiting yeast growth and extending the shelf-life of whey cheese [113].

A good temperature control is necessary for effective MAP use because the CO<sub>2</sub> effect is enhanced as the temperature decreases. A combination of refrigeration and the storage of milk and milk products under modified gas atmospheres for the extension of shelf life has been reviewed extensively [108,116–118]. In general, most of the used atmospheres do not completely inhibit growth, but the spoilage time is delayed. Overall, more research is needed to improve our knowledge on O<sub>2</sub> and CO<sub>2</sub> effects on the germination and growth of spoilage fungi and, thus, to improve MAP efficiency.

### 3.2.4. Chemical Preservatives

In addition to the methods mentioned above, chemical preservatives are widely used in the dairy industry. Preservatives are food additives that help to prevent dairy products from spoilage by bacteria and/or fungi. Antifungal preservatives used in dairy products include weak organic acids, such as sorbic acid, benzoic acid, and propionic acid, and their salts, such as potassium sorbate, calcium sorbate, sodium benzoate, potassium benzoate, calcium benzoate, and sodium propionate, as well as natamycin, a polyethylene antibiotic (Table 3).

**Table 3.** List and regulations of chemical preservatives authorized in dairy products in the European Union (EU) [119] and the United States of America (USA) [120].

Preservatives	Dairy Product	USA Regulation	Food and Drug Administration (FDA) Code	EU Legislation	EU Code
Natamycin (pimaricin)	Cheese	20 mg/kg	172.155	-	-
	Uncut hard, semi-hard, and semi-soft cheese	20 mg/kg	172.155	1 mg/dm <sup>2</sup> , surface (not present at a depth of 5 mm)	E 235
Sorbic acid	Margarine	1000 mg/kg	182.3089	-	E 200
	Flavoured fermented milk	2000 mg/kg		1500 mg/kg	
	Non-heat-treated dairy-based desserts	2000 mg/kg		300 mg/kg	
	Whey cheeses, cheese products, processed cheeses	2000 mg/kg		2000 mg/kg	
	Curdled milk, unripened products, ripened products, pre-packed, sliced; layered ripened products	2000 mg/kg		1000 mg/kg	
Potassium sorbate and calcium sorbate	Cold-pack cheese, cream cheese, pasteurized process cheese food, pasteurized process cheese spread, semi-soft part-skim cheeses	3000 mg/kg	182.3640 and 182.3225	-	E 202 and E 203
	Flavoured fermented milk products including heat-treated products	3000 mg/kg		1500 mg/kg	
	Non-heat-treated dairy-based desserts	3000 mg/kg		300 mg/kg	
	Whey cheeses, cheese products, processed cheeses	3000 mg/kg		2000 mg/kg	
	Curdled milk, unripened products, ripened products, pre-packed, sliced; layered ripened products	3000 mg/kg		1000 mg/kg	
	Flavoured fermented milk products including heat-treated products	3000 mg/kg		1500 mg/kg	
	Non-heat-treated dairy-based desserts	3000 mg/kg		300 mg/kg	
	Whey cheeses, cheese products, processed cheeses	3000 mg/kg		2000 mg/kg	
	Curdled milk, unripened products, ripened products, pre-packed, sliced; layered ripened products	3000 mg/kg		1000 mg/kg	
Sodium benzoate	Margarine	1000 mg/kg	184.1733	-	E 211
	Flavoured fermented milk products including heat-treated products	-		1500 mg/kg	
	Non-heat-treated dairy-based desserts	-		300 mg/kg	
	Whey cheeses, cheese products, processed cheeses	-		2000 mg/kg	
	Curdled milk, unripened products, ripened products, pre-packed, sliced; layered ripened products	-		1000 mg/kg	
Potassium benzoate and calcium benzoate	Flavoured fermented milk products including heat-treated products	-	184.1081	1500 mg/kg	E 212 and E 213
	Non-heat-treated dairy-based desserts	-		300 mg/kg	
	Whey cheeses, cheese products, processed cheeses	-		2000 mg/kg	
	Curdled milk, unripened products, ripened products, pre-packed, sliced; layered ripened products	-		1000 mg/kg	
Propionic acid	Gruyere cheese, swiss cheese, and emmentaler cheese	surface		-	E 280
	Ripened cheese	-		surface treatment	
Sodium propionate	Cheeses and related cheese products	surface	184.1784	-	E 281
	Ripened cheese	-		surface treatment	

Weak acids inhibit both bacterial and fungal growth. In addition, sorbic acid also inhibits spore germination in bacteria [121,122]. Preservatives such as benzoic and sorbic acid have an optimal inhibitory activity at pH between 4.5 and 5.5. Indeed, at such pH, the uncharged, undissociated acid form can diffuse freely across the cytoplasmic membrane and enter into the cell. The equilibrium between the undissociated and dissociated forms depends on the acid pKa and the medium pH. Once inside the cell (pHi ~ 7), the acid will dissociate and release charged anions and protons which cannot diffuse back across the membrane. Anions and protons accumulation into the cell is responsible for intracellular pH decrease and may cause membrane disruption, inhibition of essential metabolic reactions, stress in intracellular pH homeostasis, and/or accumulation of toxic anions which finally lead to cell death [14]. In a recent study, Garnier et al. [49] determined the minimal inhibitor concentration (MIC) of different weak acids including potassium sorbate, calcium propionate, and sodium benzoate for several fungal strains isolated from spoiled dairy products. They found that, independent of the studied species, calcium propionate was the least effective preservative followed by sodium benzoate and potassium sorbate. It should be emphasized that the inhibitory effect of weak acids on fungi is also influenced by other factors such as  $a_w$ . For example, the MIC of sorbic acid were 0.1% and 0.15% ( $w/w$ ) at 0.85  $a_w$  while they were 0.075% and 0.05% ( $w/w$ ) at 0.90  $a_w$ , for *Aspergillus flavus* and *P. roqueforti*, respectively [123]. Moreover, Stratford et al. [124] recently showed that, in contrast to acetic acid, sorbic acid did not decrease pHi in *S. cerevisiae* but instead was a membrane-active compound, inhibiting the activity of a plasma-membrane  $H^+$ -ATPase proton pump. Most studies on the action mechanism of weak acids have been performed on *S. cerevisiae* and *Zygosaccharomyces bailli*. Therefore, more studies are necessary to understand the exact inhibition mechanism of weak acids against spoilage fungi encountered in dairy products. Finally, for hard and semi-hard smear cheeses, which are characterized by the development of a red-orange microbial mat on their surfaces, films permeable to moisture and  $O_2$  can be utilized to pack the cheeses in order to protect them from yeast and molds present in the dairy environment. Such films can also be coated with antifungal molecules such as natamycin. It is worth mentioning that a successful attempt has been made to replace synthetic film polymers by a chitosan coating containing natamycin for protecting semi-hard cheese from fungal contamination [125].

Fungal resistance or adaptation to weak organic acids rely on several mechanisms. In *S. cerevisiae* and other fungi, it includes maintenance of the cell wall structure, changes in plasma membrane or cell wall composition, metal metabolism, and activation of ATP-consuming membrane transporters to remove protons and anions [126]. Ullah et al. [127] showed that *S. cerevisiae* adaptation to sorbic or acetic acid resulted in a decreased diffusional entry of the molecule. They concluded that pre-exposed cells indeed decreased acid entry through alteration of either the plasma membrane structure or the cell wall composition or through an increase in intracellular buffering capacity. In addition, Brandao et al. [38] showed that the  $H^+$  internal concentration in *S. cerevisiae* was regulated by several systems, including plasma membrane  $H^+$ -ATPase, and that ENA1p, known for its involvement in saline or alkaline stress responses and regulation of the plasma membrane potential had an important but yet to be fully understood role in the cellular response to acid [126]. This study also demonstrated that acid stress response was dependent on calcium metabolism and was blocked by a calcineurin inhibitor. Other resistance mechanisms include the metabolization of weak acids into the cell. For example, Casas et al. [128] showed that *D. hansenii* could metabolize potassium sorbate into pentadiene, a volatile compound also responsible for off-odor, while *A. niger* was able to decarboxylate and detoxify sorbic acid thanks to a phenylacrylic acid decarboxylase [129].

Natamycin (also known as pimaricin) is a fungicide belonging to polyethylene antibiotics [130]. It is currently used to control fungal growth on cheese surfaces [131–133]. Produced by *Streptomyces natalensis*, this antibiotic acts by inhibiting the vacuole fusion process through specific interaction with ergosterol at the early priming stage of fusion, but does not permeabilize the membrane.

A World Health Organization (WHO) monograph on Food Additives states that obtaining resistance against natamycin is difficult, because of the action mode of these chemical agents [134]. Indeed, natamycin binds to ergosterol, an important component of the plasma membrane and of

the growing tips of germinating spores and vegetative hyphae [135,136]. In addition, natamycin MIC of different fungal species were reported to only differ negligibly, which is not in favor of resistance development [134]. For example, *C. parapsilosis*, *Rhodotorula* spp., and *Penicillium* spp. had mean MIC of 5.2 µg/mL, 2.3 µg/mL, and 2.3 µg/mL, respectively [137]. In contrast, Garnier et al. [49], who evaluated natamycin MIC after surface-treatment of one isolate each of 10 species from spoiled dairy products, observed that MIC ranged from 0.04 mg/dm<sup>2</sup> in *Cladosporium halotolerans* to >0.2 mg/dm<sup>2</sup> in *Y. lipolytica*. In addition, a recent study [138] showed that a continuous and prolonged incubation with natamycin induced a tolerance in individual strains.

The WHO [139], European Food Safety Authority [119], and Food and Drug Administration [120] have evaluated natamycin thoroughly and all list it as safe for human consumption. In the US, according to the Code of Federal Regulations, the final amount of natamycin must not exceed 20 ppm in cheese (20 mg/kg). In the EU, natamycin is only allowed for cheese surface-treatment with a maximum applied dose of 1 mg/dm<sup>2</sup>. Moreover, it must not be present at a depth >5 mm in the finished product [140].

### 3.2.5. Fermentation

Fermentation is one of the oldest preservation methods [141,142]. It is a biological process relying on the activity of specific microorganisms producing certain metabolites with antifungal and antibacterial activities [12,141]. Different microbial groups involved in dairy product manufacturing may suppress or retard the growth of spoilage fungi. LAB produce lactic acid as a major fermentation end-product, but may also produce other metabolites with antifungal activity (see also Section 4). Certain cheese types such as smear cheeses and mold-ripened cheeses also harbor on their surfaces complex microbiota composed of desirable aerobic bacteria and fungi which can compete with spoilage fungi for one or more limiting macro- and/or micronutrients and/or for space.

A complementary approach to reduce fungal spoilage in dairy products is gaining more and more interest: the use of bioprotective cultures and fermentates.

## 4. Bioprotective Cultures

Today, in more economically developed countries, there is a strong and increasing demand from consumers for foods that are more “natural”, i.e., less heavily processed and preservative-free [75,143]. That is the reason why a strong market demand exists for natural solutions to ensure both food safety and food shelf-life [144]. Biopreservation is not a new concept as it has been used for thousands of years in fermented foods. Also called biocontrol, it refers to the extension of food shelf-life and increase in food safety using natural or added microbiota and/or their antimicrobial compounds [145]. Food bioprotective cultures can thus be defined as food-grade bacterial or fungal strains that have been selected for their antimicrobial properties. They differ from starter or adjunct cultures which are primarily used for their technological functions (acid and aroma production, role in texture, color, etc.). Among microorganisms possessing antimicrobial properties, LAB, produce a large array of antimicrobial substances including organic acids such as lactic and acetic acids, fatty acids, reuterin, antifungal peptides, and bacteriocins [146, 147]. While research has mainly focused on screening for bacterial strains with antimicrobial activity against pathogenic or spoilage bacteria and fungi, fungi with antimicrobial activities also exist [148].

### 4.1. Lactic Acid Bacteria and Propionibacteria with Antifungal Activity

LAB and propionibacteria (PAB) can be used in food production as starter cultures to modify and improve nutritional and organoleptic food properties, or as protective cultures to improve product safety and/or shelf-life [144,149]. Indeed, LAB are used as starter cultures in the manufacturing of dairy products such as fermented milk, yoghurt, buttermilk, cottage cheeses, hard cheeses, and soft cheeses, among many others [150]. LAB have also been traditionally used as natural biopreservatives of food and feed, including milk and dairy products.

Antifungal LAB have been studied in a large food range including dairy products [15,151]. Today, several recent studies report the use of antifungal LAB to control dairy product spoilage [15,152,153].



LAB produce many antifungal metabolites [154] and most LAB are granted with a generally recognized as safe (GRAS) and qualified presumption of safety (QPS) status. Besides being safe for human consumption (absence of biogenic amine production and acquired antibiotic resistance), the main properties expected from antifungal LAB and PAB bioprotective cultures are: (1) an antifungal activity exhibited and maintained during manufacturing and storage; (2) no impact on starter cultures' functionalities; (3) no modification of product organoleptic properties; (4) an activity at the lowest possible inoculum to reduce the cost associated with their use; and (5) an easy propagation at high populations and resistance to lyophilization or freezing. There are currently several antifungal bioprotective cultures commercially available for dairy products, such as HoldBac series (DuPont Danisco), FreshQ<sup>®</sup> series (Dupont), and Befresh<sup>™</sup> AF (Handary). Holdbac YM-B is a mixed culture of *Lactobacillus rhamnosus* and *Propionibacterium freudenreichii* subsp. *shermanii* and FreshQ<sup>®</sup> 2 is a single *L. rhamnosus* culture, while Befresh<sup>™</sup> AF is a mixture of *Lactobacillus paracasei* and *P. freudenreichii* subsp. *shermanii*. Despite such bioprotective cultures being available on the market, research for new LAB with antifungal activities and identification of their associated metabolites is now the focus of many academic and industrial research groups.

Screening for antifungal strains is a critical step among those required to develop antifungal cultures. For example, a recent study described a high-throughput screening method to detect antifungal activities in *Lactobacillus* species cultivated in Man, Rogosa, and Sharpe (MRS) media [155]. This method allowed the detection of 154 strains with antifungal activities against *R. mucilaginosa* and to a lower extent against *Aspergillus tamari*, *Candida krusei*, and *K. marxianus*. However, as mentioned in several papers [156,157], screening in conventional MRS medium is not recommended as MRS contains acetate which may potentialize antifungal activity and artificially inflate the number of active isolates. Another interesting but labor-consuming approach, used by Delavenne et al. [158] to develop an antifungal isolate collection, was to plate raw milk samples on eight semi-selective media for LAB, and to systematically screen colonies for their antifungal activity against four spoilage fungi using the agar-overlay method. Among the ~72,000 tested colonies, >1200 colonies (i.e., 1.7% of tested colonies) had a detectable antifungal activity. However, one should keep in mind that laboratory media differ significantly in their physicochemical and microbiological characteristics with those of dairy foods and, thus, active strains in laboratory media may lose this ability in real products. Overall, further work is needed to develop high-throughput screening methods in dairy products mimicking models to increase the chances of finding suitable strains.

Among antifungal LAB, *Lactobacillus* and, to a smaller extent, *Lactococcus*, *Pediococcus*, *Weissella*, and *Leuconostoc* are the most frequently cited genera (Table 4). Indeed, many strains pertaining to species of the *Lactobacillus* genus including *Lactobacillus plantarum*, *Lactobacillus casei*, *Lactobacillus paracasei*, and *Lactobacillus brevis* were shown to possess antifungal activity against a large spectrum of fungal targets including *Penicillium*, *Candida*, *Kluyveromyces*, and *Rhodotorula* spp. as well as *Debaryomyces hansenii* and *Yarrowia lipolytica*, which are among the most important spoilage agents in dairy products [49]. Among *Propionibacterium*, *P. freudenreichii* is the major species reported to possess antifungal activity [157,159].

So far, besides lactic, acetic, and propionic acids, which are produced at g/L or g/kg levels, a very large variety of molecules has been reported to be responsible for antifungal activity. These molecules are generally produced at lower levels (mg/L or mg/kg), and include organic acids (2-pyrrolidone-5-carboxylic acid, 3-phenyllactic acid, 4-hydroxybenzoic acid, azelaic acid, DL- $\beta$ -hydroxyphenyllactic acid, hydroxyphenyllactic, *p*-coumaric acid, phenyllactic acid (S)-(-)-2-hydrocinnamic acid, succinic acid, vanillic acid), fatty acids (3-hydroxydecanoic acid, decanoic acid, hydroxyisocaproic acid), cyclopeptides (cyclo(L-Pro-L-Pro), cyclo(L-Leu-L-Pro), cyclo(L-Tyr-L-Pro) cyclo(L-Met-L-Pro), cyclo(Phe-Pro), cyclo(Phe-OH-Pro), cyclo(L-Phe-L-Pro), cyclo(L-Phe-trans-4-OH-L-Pro) and cyclo(L-His-L-Pro)), reuterin [160], and volatile compounds such as diacetyl [153] (Table 4). Because these molecules are produced at quantities below their MIC, it is likely that they act in synergy. For example, in a study focusing on the effect of organic acids on molds, Dagnas et al. [161] showed that lactic acid alone had almost no inhibitory effect against several mold species while lactic and acetic acids could act in synergy.

**Table 4.** Lactic and propionic acid bacteria showing antifungal activities and their responsible compounds.

Lactic Acid Bacteria (LAB) and Propionibacteria	Matrix	Metabolites	Targets	References
<b>Genus <i>Lactococcus</i></b>				
<i>Lc. lactis</i>	Lab-Lemco tryptone broth (LTB)	Nd *	<i>Aspergillus flavus</i>	[162]
	LTB	Nisin	<i>Aspergillus parasiticus</i>	[163]
	Potatoe Dextrose Agar (PDA) + 0.1% Triton X-100	Nd *	<i>Aspergillus fumigatus</i> , <i>A. parasiticus</i> , <i>A. flavus</i>	[164]
	PDA + 0.1% Triton X-100	Possibly proteinaceous compound(s)	<i>Fusarium</i> spp., <i>A. flavus</i> , <i>A. parasiticus</i>	[165]
	LTB and PDA	Possibly proteinaceous compound(s), lactic acid	<i>Penicillium expansum</i>	[166]
<b>Genus <i>Lactobacillus</i></b>				
<i>Lactobacillus</i> spp.	De Man, Rogosa and Sharpe (MRS) agar	Acetic acid, propionic acid, lactic acid, peptides	<i>Penicillium candidum</i> , <i>Debaryomyces hansenii</i>	[167]
<b>Group <i>Lb. alimentarius/Lb. farciminis</i></b>				
<i>Lb. paralimentarius</i>	Modified Sabouraud Dextrose Broth (mSDB) medium	Lactic acid, phenyllactic acid, acetic acid, peptides	<i>Aspergillus japonicus</i> , <i>Eurotium repens</i> , <i>Penicillium roseopurpureum</i>	[168]
<b>Group <i>Lb. brevis</i></b>				
<i>Lb. brevis</i>	PDA	Peptide	<i>Penicillium roqueforti</i>	[169]
	wheat flour hydrolysate (WFH) broth	Acetic acid, phenyllactic acid, lactic acid	<i>Fusarium graminearum</i>	[170]
	mMRS agar	Organic acids and proteinaceous compounds	<i>Fusarium</i> spp.	[171]
	MRS agar or PDA	Peptide	<i>Penicillium camemberti</i> , <i>P. roqueforti</i> , <i>Aspergillus niger</i> , <i>Rhizopus oryzae</i> , <i>Cluyveromyces marxianus</i> , <i>Torulopsis candida</i> , <i>Meyerozyma guilliermondii</i> , <i>Saccharomyces cerevisiae</i>	[172]
<b>Group <i>Lb. casei</i></b>				
<i>Lb. casei</i>	PDA + 0.1% Triton X-100	Possibly proteinaceous compound(s)	<i>A. flavus</i> , <i>A. parasiticus</i> , <i>Fusarium</i> sp.	[165]
	PDA	Peptide	<i>Penicillium citrinum</i> , <i>P. expansum</i> , <i>A. flavus</i>	[173]
	Yoghurt	Lactic acid and cyclo-(Leu-Pro)	<i>Penicillium</i> sp.	[174]
	LTB and PDA	Possibly proteinaceous compound(s), lactic acid	<i>P. expansum</i>	[166]
<i>Lb. paracasei</i>	Yoghurt and cheese surface	Propionic acid, acetic acid, lactic acid, succinic acid, 2-pyrrolidone-5-carboxylic acid, 3-phynyllactic acid, hydroxyphenyllactic acid	<i>Candida pulcherrima</i> , <i>Candida magnoliae</i> , <i>Candida parapsilosis</i> , <i>Zygosaccharomyces bailii</i>	[151]
	MRS agar	Peptide	<i>Candida albicans</i> , <i>Candida blankii</i> , <i>Candida pseudointermedia</i>	[175]
	Chemically defined interaction medium	Diacetyl	<i>Penicillium solitum</i> , <i>Penicillium</i> sp.	[153]
	Yoghurt and acidified milk	Diacetyl, acetic acid, butanoic acid, 2,3-pentadione	<i>P. solitum</i> , <i>Penicillium</i> sp.	[153]
<i>Lb. rhamnosus</i>	Yoghurt	Acetic acid, lactic acid	<i>Rhodotorula mucilaginosa</i>	[15,158]
<b>Group <i>Lb. coryniformis</i></b>				
<i>Lb. coryniformis</i>	MRS agar or PDA	Peptide +/- 3 KDa, phenyllactic acid, cyclo(Phe-Pro), cyclo(Phe-OH-Pro), reuterin	Broad spectrum	[147]
<b>Group <i>Lb. delbrueckii</i></b>				
<i>Lb. acidophilus</i>	PDA + 0.1% Triton X	Nd *	<i>A. fumigatus</i>	[164]
<i>Lb. amylovorus</i>	MRS broth	3-phenylpropanoic acid, <i>p</i> -coumaric, (E)-2-methylcinnamic acid, 3-phenyllactic acid, 3-(4-hydroxyphenyl)lactic acid, lactic acid, acetic acid, D-glucuronic acid, salicylic acid, cytidine and 2'-deoxycytidine, sodium decanoate,	<i>A. fumigatus</i> , <i>Fusarium culmorum</i>	[175]

Table 4. Cont.

Lactic Acid Bacteria (LAB) and Propionibacteria	Matrix	Metabolites	Targets	References
	Cheese	Cyclo(L-Pro-L-Pro), cyclo(L-Leu-L-Pro), cyclo(L-Tyr-L-Pro) cyclo(L-Met-L-Pro) and cyclo(L-His-L-Pro)	<i>P. roqueforti</i> , <i>P. expansum</i>	[176]
	Milk agar and cheese	DL- <i>p</i> -hydroxyphenyllactic acid, 4-hydroxybenzoic acid, (S)-(–)-2-hydroxyisocaproic acid, azelaic acid, phenyllactic acid, benzoic acid, hydrocinnamic acid, 3-hydroxydecanoic acid, DL-β-hydroxylauric acid, decanoic acid, salicylic acid, 4-hydroxybenzoic, vanillic acid, (S)-(–)-2-hydroxyisocaproic acid	<i>P. expansum</i>	[177]
<i>Lb. delbrueckii</i>	LTB and PDA	Possibly proteinaceous compound(s), lactic acid	<i>P. expansum</i>	[166]
Group <i>Lb. fructivorans</i>				
<i>Lb. sanfranciscensis</i>	Malt-agar medium	Caproic acid, propionic acid, butyric acid, acetic acid, valeric acid	<i>F. graminearum</i>	[178,179]
Group <i>Lb. perolens</i>				
<i>Lb. harbinensis</i>	Yoghurt	Acetic acid, lactic acid	<i>Yarrowia lipolytica</i> , <i>P. expansum</i> , <i>Penicillium brevicompactum</i> , <i>D. hansenii</i> , <i>R. mucilaginosa</i> , <i>Kluyveromyces lactis</i>	[15,158]
Group <i>Lb. plantarum</i>				
<i>Lb. pentosus</i>	MRS	Peptide	<i>A. niger</i>	[180]
	MRS agar	Peptide, phenyllactic and hydroxyphenyllactic acid	<i>Penicillium nalgiovense</i> , <i>Aspergillus candidus</i>	[181]
<i>Lb. plantarum</i>	mMRS agar	(S)-(–)-2-hydroxyisocaproic acid, hydrocinnamic acid, phenyllactic acid, decanoic acid, azelaic acid, 4-hydroxybenzoic acid, <i>p</i> -coumaric acid, vanillic acid, DL- <i>p</i> -hydroxyphenyllactic acid, 3-hydroxydecanoic acid	<i>Microsporium canis</i> , <i>Microsporium gypseum</i> , <i>Epidermophyton floccosum</i>	[182]
	MRS broth	Benzoic acid, 5-methyl-2,4-imidazolidinedione, tetrahydro-4-hydroxy-4-methyl-2H-pyran-2-one, 3-(2-methylpropyl)-2,5-piperazinedione, cyclo(glycyl-L-leucyl)).	<i>Fusarium avenaceum</i>	[183]
	MRS agar	Lactic acid, PLA, cyclo(L-Leu-L-Pro), cyclo(L-Phe-L-Pro)	<i>A. niger</i> , <i>F. graminearum</i> , <i>F. culmorum</i> , <i>Fusarium oxysporum</i>	[184]
	Wheat flour hydrolysate (WFH)	Phenyllactic acid, 4-hydroxy-phenillactic acid	<i>E. repens</i> , <i>Eurotium rubrum</i> , <i>Penicillium corylophilum</i> , <i>P. roqueforti</i> , <i>P. expansum</i> , <i>Endomyces fibuliger</i> , <i>A. niger</i> , <i>A. flavus</i> , <i>Monilia sitophila</i> , <i>F. graminearum</i>	[185]
	MRS broth	3-phenyllactic acid, cyclo(Phe-Pro), cyclo(Phe-OH-Pro), cyclo(L-Phe-LPro) and cyclo(L-Phe-trans-4-OH-L-Pro) dipeptides	<i>Fusarium sporotrichioides</i> , <i>A. fumigatus</i> , <i>K. marxianus</i>	[186]
	Wheat flour hydrolysate (WFH)	Acetic acid, phenyllactic acid, lactic acid	<i>F. graminearum</i> , <i>A. niger</i>	[170]
	MRS agar	3-( <i>R</i> )-hydroxydecanoic acid, 3-hydroxy-5-cis-dodecanoic acid, 3-( <i>R</i> )-hydroxydodecanoic acid and 3-( <i>R</i> )-hydroxytetradecanoic acid	<i>A. fumigatus</i> , <i>Aspergillus nidulans</i> , <i>K. marxianus</i> , <i>P. roqueforti</i> , <i>Penicillium commune</i> , <i>Penicillium anomala</i> , <i>R. mucilaginosa</i>	[187]
	MRS agar plates	Acetic acid	<i>A. flavus</i> , <i>F. graminearum</i> , <i>Rhizopus stolonifer</i> , <i>Sclerotium oryzae</i> , <i>Rhizoctonia solani</i> , <i>Botrytis cinerea</i> , <i>Sclerotinia minor</i>	[188]
	MRS agar	Peptide, phenyl-lactic and hydroxy-phenyllactic acid	<i>P. nalgiovense</i> , <i>P. camemberti</i> , <i>Penicillium verrucosum</i> , <i>Penicillium chrysogenum</i> , <i>A. candidus</i> , <i>A. flavus</i> , <i>A. ochraceus</i> , <i>A. fumigatus</i> , <i>Galactomyces geotrichum</i> , <i>Moniliella</i> spp., <i>Mucor racemosus</i> , <i>Wallemia sebi</i> , <i>Eurotium herbariorum</i>	[181]
	Soybean	3,6-bis(2-methylpropyl)-2,5-piperazinedion	<i>A. flavus</i>	[189]
	MRS agar plates	3-3-phenyllactic acid (PLA), lactic acid, acetic acid	<i>A. fumigatus</i> , <i>Rhizopus stolonifer</i>	[190]

Table 4. Cont.

Lactic Acid Bacteria (LAB) and Propionibacteria	Matrix	Metabolites	Targets	References
	MRS agar medium, apple-based agar growth medium	Lactic acid, acetic acid	<i>P. expansum</i> , <i>Penicillium notatum</i>	[191]
	MRS agar medium	2-hydroxy-4-methylpentanoic acid	<i>A. niger</i> , <i>Aspergillus tubingensis</i> , <i>Penicillium crustosum</i>	[192]
	PDA plates	3-PLA, benzeneacetic acid, 2-propenyl ester	<i>B. cinerea</i> , <i>Glomerella cingulate</i> , <i>Phytophthora drechsleri</i> Tucker, <i>P. citrinum</i> , <i>Penicillium digitatum</i> , <i>F. oxysporum</i>	[193]
	Chopped Meat Carbohydrate (CMC) broth	Lactic acid	<i>F. avenaceum</i> , <i>F. culmorum</i> , <i>F. graminearum</i> , <i>F. oxysporum</i>	[194]
<b>Group <i>Lb. reuteri</i></b>				
<i>Lb. fermentum</i>	MRS	Peptide	<i>A. niger</i>	[180]
<i>Lb. reuteri</i>	Wheat flour hydrolysate (WFH) broth	Acetic acid, phenyllactic acid, lactic acid	<i>F. graminearum</i> , <i>A. niger</i>	[170]
	mMRS agar	Acid, vanillic acid, DL-P-hydroxyphenyllactic acid, 3-hydroxydecanoic acid, (S)-(-)-2-hydroxyisocaproic acid, hydrocinnamic acid, phenyllactic acid, decanoic acid, azelaic acid, 4-hydroxybenzoic acid, p-coumaric	<i>M. canis</i> , <i>M. gypseum</i> , <i>E. floccosum</i>	[182]
<i>Lb. rossiae</i>	mSDB agar medium	Lactic acid, phenyllactic acid, acetic acid, peptides	<i>A. japonicus</i> , <i>E. repens</i> , <i>P. roseopurpureum</i>	[168]
<b>Group <i>Lb. sakei</i></b>				
<i>Lb. sakei</i>	MRS	Sakacin KTU05-6	<i>A. flavus</i> , <i>A. fumigatus</i> , <i>A. niger</i> , <i>Aspergillus versicolor</i> , <i>F. culmorum</i> , <i>Fusarium poae</i> , <i>Mucor</i> spp., <i>P. chrysogenum</i> , <i>P. expansum</i> , <i>Penicillium</i> spp.	[195]
<b>Genus <i>Pediococcus</i></b>				
<i>Pc. acidilactici</i>	MRS	Pediocin KTU05-08	<i>A. flavus</i> , <i>A. fumigatus</i> , <i>A. niger</i> , <i>A. versicolor</i> , <i>F. culmorum</i> , <i>F. poae</i> , <i>Mucor</i> spp., <i>P. chrysogenum</i> , <i>P. expansum</i> , <i>Penicillium</i> spp.	[195]
<i>Pc. pentosaceus</i>	MRS	Peptide	<i>A. niger</i>	[180]
	MRS agar plates	Acetic acid	<i>A. flavus</i> , <i>F. graminearum</i> , <i>R. stolonifer</i> , <i>S. oryzae</i> , <i>R. solani</i> , <i>B. cinerea</i> , <i>S. minor</i>	[188]
	MRS	Pediocin KTU05-09, KTU05-10 and AcKTU05-67	<i>A. flavus</i> , <i>A. fumigatus</i> , <i>A. niger</i> , <i>A. versicolor</i> , <i>F. culmorum</i> , <i>F. poae</i> , <i>Mucor</i> spp., <i>P. chrysogenum</i> , <i>P. expansum</i> , <i>Penicillium</i> spp.	[195]
	MRS agar medium	Possibly cyclic dipeptide	<i>P. expansum</i> , <i>Penicillium notatum</i>	[191]
<b>Genus <i>Weissella</i></b>				
<i>W. confusa</i>	MRS agar medium	Lactic acid, acetic acid	<i>P. expansum</i> , <i>P. notatum</i>	[191]
<i>W. cibaria</i>	MRS agar medium	Lactic acid, acetic acid	<i>P. expansum</i> , <i>P. notatum</i>	[191]
<i>W. paramesenteroides</i>	MRS agar medium	2-hydroxy-4-methylpentanoic acid, lactic acid, acetic acid	<i>A. niger</i> , <i>A. tubingensis</i> , <i>P. crustosum</i>	[192]
	MRS agar plates	Acetic acid	<i>A. flavus</i> , <i>F. graminearum</i> , <i>R. stolonifer</i> , <i>S. oryzae</i> , <i>B. cinerea</i> , <i>S. minor</i>	[188]

Table 4. Cont.

Lactic Acid Bacteria (LAB) and Propionibacteria	Matrix	Metabolites	Targets	References
<b>Genus <i>Propionibacteria</i></b>				
<i>P. acidipropionici</i>	Sodium lactate (SL) medium	Lactic acid, propionic acid, acetic acid	<i>A. fumigatus</i>	[196]
	MRS	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
	MRS-acetate	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
	SL broth	PLA	<i>A. fumigatus</i>	[196]
<i>P. freudenreichii</i> subsp. <i>shermanii</i>	Skim milk	propionic acid, 3-phenyllactic acid	<i>P. chrysogenum</i>	[159]
	MRS	Acetic acid, propionic acid, 3-phenyllactic acid, 4-hydroxy-phenyllactic acid	<i>P. chrysogenum</i>	[159]
	MRS	Propionic acid, acetic acid	<i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
	MRS-acetate	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
<i>P. freudenreichii</i> subsp. <i>freudenreichii</i>	MRS	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
	MRS-acetate	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
<i>P. jensenii</i>	Yoghurt and cheese surface	Propionic acid, acetic acid, lactic acid, succinic acid, 2-pyrrolidone-5-carboxylic acid, 3-phenyllactic acid, hydroxyphenyllactic acid	<i>C. pulcherrima</i> , <i>C. magnoliae</i> , <i>C. parapsilosis</i> , <i>Z. bailii</i>	[151]
	MRS	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
	MRS-acetate	Nd *	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]
<i>P. thoenii</i>	Sodium Lactate (SL) medium	Lactic acid, propionic acid	<i>A. fumigatus</i>	[196]
	SL broth	PLA	<i>A. fumigatus</i>	[196]
	MRS	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i> , <i>K. marxianus</i>	[196]
	MRS-acetate	Propionic acid, acetic acid	<i>R. mucilaginoso</i> , <i>P. roqueforti</i> , <i>A. fumigatus</i>	[196]

\* Nd: not determined.



Similar results were reported for other organic acids such as PLA, in which MIC decreased in the presence of lactic acid [185]. Only few studies [154,197] dealt with the action mode of LAB antifungal metabolites and the response of fungi at the physiological, transcriptomic, or proteomic levels. Therefore, further research should be undertaken in this area.

As shown in Table 3, most studies on LAB and PAB antifungal activity have been performed in (semi-)synthetic culture medium and it is obvious that important discrepancies can exist between in vitro and in situ tests. The main reasons for this are that intrinsic and extrinsic factors can affect the production of antifungal metabolites, their activity, and the susceptibility of the fungal target to these compounds. Intrinsic factors include medium composition in terms of macro- and micronutrients, pH,  $a_w$ , Eh, and food structure, while extrinsic factors are temperature, composition of the surrounding atmosphere, and humidity. Therefore, a second critical step in developing antifungal cultures is to test their efficiency in real products using challenge- and durability-tests and against one or several fungal targets. It should be noted that only a few publications have clearly shown in situ antifungal activity of selected LAB and PAB (Table 4). For example, Delavenne et al. [15,27] showed the antifungal activity of *Lactobacillus harbinensis* KV931Np against six fungal targets in yoghurt while Schwenninger and Meile (2004) demonstrated the antifungal activity of three mixed cultures of *Lactobacillus paracasei* subsp. *paracasei* SM20, SM29, or SM63 and *Propionibacterium jensenii* SM11, against *Candida pulcherrima*, *C. magnoliae*, *C. parapsilosis*, and *Zygosaccharomyces bailii* in yoghurt and on cheese surfaces [151]. More recently, Aunsbjerg et al. [153] proved the antifungal activity of *Lactobacillus paracasei* in chemically defined medium and in yoghurt against *P. solitum* and *Penicillium* sp.

#### 4.2. Fermentates

LAB and PAB may also be used to produce dairy fermentates containing antifungal metabolites, which are fermented dairy ingredients produced from milk via a fermentation process [144]. MicroGARD (DuPont Danisco) and DuraFresh (Kerry) are two currently available commercial fermentates. As an example, MicroGARD, which is FDA-approved, is produced by skimmed milk fermentation using *P. freudenreichii* subsp. *shermanii*. The use of this fermentate in cottage cheese partially inhibited *K. marxianus* and *P. expansum* and thus extended its shelf-life [197]. It may also be used in sour cream, yoghurt, and dairy desserts [198].

### 5. Conclusions

Despite technological advances, fungal spoilage is still a main issue in the dairy industry. Among the actual methods in use, a large focus concerns the replacement of traditional hurdle technologies such as chemical preservatives by new techniques to meet the increasing consumer demand for less-heavily processed and preservative-free dairy products. These new techniques include preventive methods such as a better management of air quality and non-thermal control methods such as modified atmosphere packaging and biopreservation using antifungal bioprotective cultures or fermentates.

**Acknowledgments:** We are grateful to the regional councils of Brittany (grant No. 13008651) and Pays de la Loire (grant No. 2014-07081) for the financial support and INRA for scientific coordination (J. Leonil) through the interregional project PROFIL, managed by BBA industrial association.

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

1. FAO. Global FOOD Losses and Food Waste—Extent, Causes and Prevention. Rome. 2011. Available online: <http://www.fao.org/docrep/014/mb060e/mb060e00.pdf> (accessed on 5 July 2017).
2. Pitt, J.I.; Hocking, A.D. *Fungi and Food Spoilage*, 3rd ed.; Springer Science & Business Media: New York, NY, USA, 2009; pp. 1–2, ISBN 978-1-4615-6391-4.
3. Shephard, G.S. Impact of mycotoxins on human health in developing countries. *Food Addit. Contam. A* **2008**, *25*, 146–151. [CrossRef] [PubMed]

4. Filtenborg, O.; Frisvad, J.C.; Thrane, U. Moulds in food spoilage. *Int. J. Food Microbiol.* **1996**, *33*, 85–102. [[CrossRef](#)]
5. Huis in't Veld, J.H.J. Microbial and biochemical spoilage of foods: An overview. *Int. J. Food Microbiol.* **1996**, *33*, 1–18. [[CrossRef](#)]
6. Prado, M.R.; Blandón, L.M.; Vandenberghe, L.P.S.; Rodrigues, C.; Castro, G.R.; Thomas-Soccol, V.; Soccol, C.R. Milk kefir: Composition, microbial cultures, biological activities and related products. *Front. Microbiol.* **2015**, *6*, 1177. [[CrossRef](#)] [[PubMed](#)]
7. Desmaures, N. CHEESE-mold-ripened varieties. In *Encyclopedia of Food Microbiology*, 2nd ed.; Batt, C.A., Ed.; Elsevier Inc.: Atlanta, GA, USA, 2014; pp. 409–415, ISBN 9780123847331.
8. Baroiller, C.; Schmidt, J.L. Contribution à l'étude de l'origine des levures du fromage de Camembert. *Lait* **1990**, *70*, 67–84. [[CrossRef](#)]
9. Ledenbach, L.H.; Marshall, R.T. Microbiological spoilage of dairy products. In *Compendium of the Microbiological Spoilage of Foods and Beverages, Food Microbiology and Food Safety*; Sperber, W.H., Doyle, M.P., Eds.; Springer: New York, NY, USA, 2010; pp. 41–67, ISBN 1441908269, 9781441908261.
10. Hymery, N.; Vasseur, V.; Coton, M.; Mounier, J.; Jany, J.L.; Barbier, G.; Coton, E. Filamentous fungi and mycotoxins in cheese: A review. *Compr. Rev. Food Sci. Food Saf.* **2014**, *13*, 437–456. [[CrossRef](#)]
11. Westall, S.; Filtenborg, O. Spoilage yeasts of decorated soft cheese packed in modified atmosphere. *Food Control* **1998**, *21*, 136–142. [[CrossRef](#)]
12. Ross, R.P.; Morgan, S.; Hill, C. Preservation and fermentation: Past, present and future. *Int. J. Food Microbiol.* **2002**, *79*, 3–16. [[CrossRef](#)]
13. Stratford, M.; Steels, H.; Nebe-von-Caron, G.; Avery, S.V.; Novodvorska, M.; Archer, D.B. Population heterogeneity and dynamics in starter culture and lag phase adaptation of the spoilage yeast *Zygosaccharomyces bailii* to weak acid preservatives. *Int. J. Food Microbiol.* **2014**, *181*, 40–47. [[CrossRef](#)] [[PubMed](#)]
14. Brul, S.; Coote, P. Preservatives agents in foods: Mode of action and microbial resistance mechanisms. *Int. J. Food Microbiol.* **1999**, *50*, 1–17. [[CrossRef](#)]
15. Delavenne, E.; Cliquet, S.; Trunet, C.; Barbier, G.; Mounier, J.; Le Blay, G. Characterization of the antifungal activity of *Lactobacillus harbinensis* K.V9.3.1.Np and *Lactobacillus rhamnosus* K. C8.3.1I in yoghurt. *Food Microbiol.* **2015**, *45*, 10–17. [[CrossRef](#)] [[PubMed](#)]
16. Schoch, C.L.; Seifert, K.A.; Huhndorf, S.; Robert, V.; Spouge, J.L.; Levesque, C.A.; Chen, W. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 6241–6246. [[CrossRef](#)] [[PubMed](#)]
17. Almeida, M.; Hébert, A.; Abraham, A.L.; Rasmussen, S.; Monnet, C.; Pons, N.; Delbès, S.; Loux, V.; Batta, J.M.; Leonard, P.; et al. Construction of a dairy microbial genome catalog opens new perspectives for the metagenomic analysis of dairy fermented products. *BMC Genom.* **2014**, *15*, 1257–1287. [[CrossRef](#)] [[PubMed](#)]
18. Callon, C.; Rétureau, E.; Didienne, R.; Montel, M.-C. Microbial biodiversity in cheese consortia and comparative *Listeria* growth on surfaces of uncooked pressed cheeses. *Int. J. Food Microbiol.* **2014**, *174*, 98–109. [[CrossRef](#)] [[PubMed](#)]
19. Montel, M.-C.; Buchin, S.; Mallet, A.; Delbes-Paus, C.; Vuitton, D.A.; Desmaures, N.; Berthier, F. Traditional cheeses: Rich and diverse microbiota with associated benefits. *Int. J. Food Microbiol.* **2014**, *177*, 136–154. [[CrossRef](#)] [[PubMed](#)]
20. Wolfe, B.E.; Button, J.E.; Santarelli, M.; Dutton, R.J. Cheese rind communities provide tractable systems for in situ and in vitro studies of microbial diversity. *Cell* **2014**, *158*, 422–433. [[CrossRef](#)] [[PubMed](#)]
21. Quigley, L.; O'Sullivan, O.; Stanton, C.; Beresford, T.P.; Ross, R.P.; Fitzgerald, G.F.; Cotter, P.D. The complex microbiota of raw milk. *FEMS Microbiol. Rev.* **2013**, *37*, 664–698. [[CrossRef](#)] [[PubMed](#)]
22. Dugat-Bony, E.; Garnier, L.; Denonfoux, J.; Ferreira, S.; Sarthou, A.S.; Bonnarme, P.; Irlinger, F. Highlighting the microbial diversity of 12 French cheese varieties. *Int. J. Food Microbiol.* **2016**, *238*, 265–273. [[CrossRef](#)] [[PubMed](#)]
23. Lavoie, K.; Touchette, M.; St-Gelais, D.; Labrie, S. Characterization of the fungal microflora in raw milk and specialty cheeses of the province of Quebec. *Dairy Sci. Technol.* **2012**, *92*, 455–468. [[CrossRef](#)] [[PubMed](#)]
24. Marín, P.; Palmero, D.; Jurado, M. Occurrence of moulds associated with ovine raw milk and cheeses of the Spanish region of Castilla La Mancha. *Int. J. Dairy Technol.* **2015**, *68*, 565–572. [[CrossRef](#)]

25. Vacheyrou, M.; Normand, A.C.; Guyot, P.; Cassagne, C.; Piarrou, R.; Bouton, Y. Cultivable microbial communities in raw cow milk and potential transfers from stables of sixteen French farms. *Int. J. Food Microbiol.* **2011**, *146*, 253–262. [[CrossRef](#)] [[PubMed](#)]
26. Fleet, G.H. Yeasts in dairy products. *J. Appl. Bacteriol.* **1990**, *68*, 199–211. [[CrossRef](#)] [[PubMed](#)]
27. Von Neubeck, M.; Baur, C.; Krewinkel, M.; Stoeckel, M.; Kranz, B.; Stressler, T.; Fisher, L.; Hinrichs, J.; Scherer, S.; Wenning, M. Biodiversity of refrigerated raw milk microbiota and their enzymatic spoilage potential. *Int. J. Food Microbiol.* **2015**, *211*, 57–65. [[CrossRef](#)] [[PubMed](#)]
28. Kure, C.F.; Skaar, I.; Brendehaug, J. Mould contamination in production of semi-hard cheese. *Int. J. Food Microbiol.* **2004**, *93*, 41–49. [[CrossRef](#)] [[PubMed](#)]
29. Beletsiotis, E.; Guikas, D.; Kalantzi, K. Incorporation of microbiological and molecular methods in HACCP monitoring scheme of molds and yeasts in a Greek dairy plant: A case study. *Proc. Food Sci.* **2011**, *1*, 1051–1059. [[CrossRef](#)]
30. Temelli, S.; Anar, S.; Sen, C.; Akyuva, P. Determination of microbiological contamination sources during Turkish white cheese production. *Food Control* **2006**, *17*, 856–861. [[CrossRef](#)]
31. Oyugi, E.; Buys, E.M. Microbiological quality of shredded Cheddar cheese packaged in modified atmospheres. *Int. J. Dairy Technol.* **2007**, *60*, 89–95. [[CrossRef](#)]
32. Bokulich, N.A.; Mills, D.A. Facility-specific “house” microbiome drives microbial landscape of artisan cheesemaking plants. *Appl. Environ. Microbiol.* **2013**, *79*, 5214–5223. [[CrossRef](#)] [[PubMed](#)]
33. Radha, K.; Nath, L.S. Studies on the air quality in a dairy processing plant. *Ind. J. Vet. Anim. Sci. Res.* **2014**, *43*, 346–353. [[CrossRef](#)]
34. Schön, K.; Schornsteiner, E.; Dzieciol, M.; Wagner, M.; Müller, M.; Schmitz-Esser, S. Microbial communities in dairy processing environment floor-drains are dominated by product-associated bacteria and yeasts. *Food Control* **2016**, *70*, 210–215. [[CrossRef](#)]
35. Kaminarides, S.E.; Lakos, N.S. Yeasts in factory brine of feta cheese. *Aust. J. Dairy Technol.* **1992**, *47*, 68–71.
36. Suriyarachchi, V.R.; Fleet, G.H. Occurrence and growth of yeasts in yoghurts. *Appl. Environ. Microbiol.* **1981**, *42*, 574–579. [[CrossRef](#)] [[PubMed](#)]
37. Mayoral, M.B.; Martin, R.; Sanz, A.; Hernandez, P.E.; Gonzalez, I.; Garcia, T. Detection of *Kluyveromyces marxianus* and other spoilage yeasts in yoghurt using a PCR-culture technique. *Int. J. Food. Microbiol.* **2005**, *105*, 27–34. [[CrossRef](#)] [[PubMed](#)]
38. Penney, V.; Henderson, G.; Blum, C.; Johnson-Green, P. The potential phytopreservatives and nisin to control microbial spoilage of minimally processed fruit yoghurts. *Innov. Food Sci. Emerg. Technol.* **2004**, *5*, 369–375. [[CrossRef](#)]
39. Tranquillini, R.; Scaramuzza, N.; Berni, E. Occurrence and ecological distribution of heat resistant moulds spores (HRMS) in raw materials used by food industry and thermal characterization of two *Talaromyces* isolates. *Int. J. Food Microbiol.* **2017**, *242*, 116–123. [[CrossRef](#)] [[PubMed](#)]
40. Catellani, P.; Miotti Scapin, R.; Alberghini, L.; Radu, I.L.; Giaccone, V. Levels of microbial contamination of domestic refrigerators in Italy. *Food Control* **2014**, *42*, 257–262. [[CrossRef](#)]
41. Carreira, A.; Ferreira, L.M.; Loureiro, V. Brown pigments produced by *Yarrowia lipolytica* result from extracellular accumulation of homogentisic acid. *Appl. Environ. Microbiol.* **2001**, *67*, 3463–3468. [[CrossRef](#)] [[PubMed](#)]
42. Carreira, A.; Ferreira, L.M.; Loureiro, V. Production of brown tyrosine pigments by the yeast *Yarrowia lipolytica*. *J. Appl. Microbiol.* **2001**, *90*, 372–379. [[CrossRef](#)] [[PubMed](#)]
43. Dalgaard, P.; Gram, L.; Huss, H.H. Spoilage and shelf life of cod fillets packed in vacuum or modified atmospheres. *Int. J. Food Microbiol.* **1993**, *19*, 283–294. [[CrossRef](#)]
44. Gardini, F.; Tofalo, R.; Belletti, N.; Iucci, L.; Suzzi, G.; Torriani, S.; Guerzoni, M.E.; Lanciotti, R. Characterization of yeasts involved in the ripening of Pecorino Crotonese cheese. *Food Microbiol.* **2006**, *23*, 641–648. [[CrossRef](#)] [[PubMed](#)]
45. Roig-Sagués, A.X.; Molina, A.P.; Hernández-Herrero, M. Histamine and tyramine-forming microorganisms in Spanish traditional cheeses. *Eur. Res. Technol.* **2002**, *215*, 69–100. [[CrossRef](#)]
46. Suzzi, G.; Schirone, M.; Martuscelli, M.; Gatti, M.; Fornasari, M.E.; Neviani, E. Yeasts associated with Manteca. *FEMS Yeast Res.* **2003**, *3*, 159–166. [[CrossRef](#)]
47. Linares, D.M.; Martín, M.C.; Ladero, V.; Álvarez, M.A.; Fernández, M. Biogenic amines in dairy products. *Crit. Rev. Food Sci. Nutr.* **2011**, *51*, 691–703. [[CrossRef](#)] [[PubMed](#)]

48. Leuschner, R.G.; Heidel, M.; Hammes, W.P. Histamine and tyramine degradation by food fermenting microorganisms. *Int. J. Food Microbiol.* **1998**, *39*, 1–10. [[CrossRef](#)]
49. Garnier, L.; Valence, F.; Pawtowski, A.; Auhustsinava-Galerie, L.; Frotté, N.; Baroncelli, R.; Deniel, F.; Coton, E.; Mounier, J. Diversity of spoilage fungi associated with French dairy products. *Int. J. Food Microbiol.* **2017**, *241*, 191–197. [[CrossRef](#)] [[PubMed](#)]
50. Robinson, R.K. Dairy microbiology. In *The Microbiology of Milk*, 2nd ed.; Literary Cat Books: London, UK; New York, NY, USA, 1990; Volume 1, ISBN 978-0-471-38596-7.
51. Minervini, F.; Montagna, M.T.; Spilotros, G.; Monaci, L.; Santacroce, M.P.; Visconti, A. Survey on mycoflora of cow and buffalo dairy products from Southern Italy. *Int. J. Food Microbiol.* **2001**, *69*, 141–146. [[CrossRef](#)]
52. Jakobsen, M.; Narchus, J. Yeasts and their possible beneficial and negative effects on the quality of dairy products. *Int. Dairy J.* **1996**, *6*, 755–768. [[CrossRef](#)]
53. Fleet, G.H.; Mian, M.A. The occurrence and growth of yeasts in dairy products. *Int. J. Food Microbiol.* **1987**, *4*, 145–155. [[CrossRef](#)]
54. Hocking, A.D.; Faedo, M. Fungi causing thread mould spoilage of vacuum packaged Cheddar cheese during maturation. *Int. J. Food Microbiol.* **1992**, *16*, 123–130. [[CrossRef](#)]
55. Fadda, M.E.; Cosentino, S.; Deplano, M.; Palmas, F. Yeasts populations in Sardinian feta cheese. *Int. J. Food Microbiol.* **2001**, *69*, 153–156. [[CrossRef](#)]
56. Montagna, M.T.; Santacroce, M.P.; Spilotros, G.; Napoli, C.; Minervini, F.; Papa, A.; Dragoni, I. Investigation of fungal contamination in sheep and goat cheeses in southern Italy. *Mycopathologia* **2004**, *158*, 245–249. [[CrossRef](#)] [[PubMed](#)]
57. Jodral, M.; Liñan, E.; Acosta, I.; Gallego, C.; Roja, F.; Bentabol, A. Mycoflora and toxigenic *Aspergillus flavus* in Spanish milks. *Int. J. Food Microbiol.* **1993**, *36*, 171–174. [[CrossRef](#)]
58. Sensidoni, A.; Rondinini, G.; Peressini, D.; maifreni, M.; Bortolomeazzi, R. Presence of an off-flavor associated with the use of sorbates in cheese and margarine. *Int. J. Food Sci.* **1994**, *6*, 237–242.
59. Lund, F.; Filtenborg, O.; Frisvad, J.C. Associated mycoflora of cheese. *Food Microbiol.* **1995**, *12*, 173–180. [[CrossRef](#)]
60. Houbaken, J.; Samson, R.A. Phylogeny of *Penicillium* and the segregation of Trichocomaceae into three families. *Stud. Mycol.* **2011**, *70*, 1–51. [[CrossRef](#)] [[PubMed](#)]
61. Frisvad, J.C.; Houbaken, J.; Popma, S.; Samson, R.A. Two new *Penicillium* species *Penicillium buchwaldii* and *Penicillium spathulatum*, producing the anticancer compound asperphenamate. *FEMS Microbiol. Lett.* **2013**, *339*, 77–92. [[CrossRef](#)] [[PubMed](#)]
62. Samson, R.; Frisvad, J.; Hoekstra, E.; Filtenborg, O. Introduction to food and airborne fungi. In *Mycopathologia*, 6th ed.; Centraal Bureau Voor Schimmelcultures: Utrecht, The Netherlands, 2002; p. 609, ISBN 10 9070351528.
63. Moubasher, A.H.; Abdel-Kader, M.I.A.; El-Kady, I.A. Toxigenic fungi isolated from Roquefort cheese. *Mycopathologia* **1978**, *66*, 187–190. [[CrossRef](#)]
64. Kikoku, Y.; Tagashira, N.; Nakano, H. Heat resistance of fungi isolated from frozen blueberries. *J. Food Prot.* **2008**, *71*, 2030–2035. [[CrossRef](#)] [[PubMed](#)]
65. Kure, C.F.; Wasteson, Y.; Brendehaug, J.; Skaar, I. Mould contaminants on Jarlsberg and Norvegia cheese blocks from four factories. *Int. J. Food Microbiol.* **2001**, *70*, 21–27. [[CrossRef](#)]
66. Bullerman, L.B.; Olivigni, L.B. Mycotoxin producing-potential of molds isolated from cheddar cheese. *J. Food Sci.* **1974**, *39*, 1166–1168. [[CrossRef](#)]
67. Hassanin, I.N. Detection of mycotoxigenic fungi and bacteria in processed cheese in Egypt. *Int. Biodeterior. Biodegrad.* **1993**, *31*, 15–23. [[CrossRef](#)]
68. Aran, N.; Eke, D. Mould mycoflora of Kaşar cheese at the stage of consumption. *Food Microbiol.* **1987**, *4*, 101–104. [[CrossRef](#)]
69. Kure, C.F.; Skaar, I. Mould growth on Norwegian semi-hard cheeses Norvegia and Jarlsberg. *Int. J. Food Microbiol.* **2000**, *62*, 133–137. [[CrossRef](#)]
70. Basílico, J.C.; deBasílico, M.Z.; Chiericatti, C.; Vinderola, C.G. Characterizaion and control of thread mould in cheese. *Lett. Appl. Microbiol.* **2001**, *32*, 419–423. [[CrossRef](#)] [[PubMed](#)]
71. Gould, G.W. *Mechanisms of Action of Food Preservation Procedures*; Gould, G.W., Ed.; Elsevier Applied Science: London, UK, 1989; ISBN 1851662936, 9781851662937.
72. Rahman, M.S. *Handbook of Food Preservation*, 2nd ed.; CRC Press: Boca Raton, FL, USA, 2007; ISBN 9781574446067.



73. Sofos, J.N. Current microbiological considerations in food preservation. *Int. J. Food Microbiol.* **1993**, *19*, 87–108. [CrossRef]
74. Dalié, D.K.D.; Deschamps, A.M.; Richard-Forget, F. Lactic acid bacteria—Potential for control of mould growth and mycotoxins: A review. *Food Control* **2010**, *21*, 370–380. [CrossRef]
75. Regulation (EC) No. 852/2004 of the European Parliament and of the Council of 29 April 2004 on the Hygiene and Foodstuffs. Available online: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:139:0001:0054:en:PDF> (accessed on 27 July 2017).
76. Rysstad, G.; Kolstad, J. Extended shelf life milk—Advances in technology. *Int. J. Dairy Technol.* **2006**, *85*–96. [CrossRef]
77. Loureiro, V.; Malfeito-Ferreira, M. Spoilage yeasts in foods and beverages: Characterization and ecology for improved diagnosis and control. *Food Res. Int.* **2000**, *33*, 247–256. [CrossRef]
78. Loureiro, V.; Querol, A. The prevalence and control of spoilage yeasts in foods and beverages. *Trends Food Sci. Technol.* **1999**, *10*, 356–365. [CrossRef]
79. Odet, G. Qualité bactériologique des fromages au lait cru. *Cahiers de Nutrition et de Diététique* **1999**, *34*, 47–53.
80. Faye, B.; Loiseau, G. *Sources of Contamination in Dairy Supply Chains and Approaches to Quality Control*; Paper Read at International Cooperation in International Research for Agricultural Development (CIRAD)-FAO: Montpellier, France, 2002; ISBN 92-5-004787-8.
81. Sandrou, D.K.; Arvanitoyannis, I.S. Implementation of Hazard Analysis Critical Control Point (HACCP) to the dairy industry: Current status and perspectives. *Food Rev. Int.* **2000**, *16*, 77–111. [CrossRef]
82. De Steur, H.; Wesana, J.; Dora, M.K.; Pearce, D.; Gellynck, X. Applying value stream mapping to reduce food losses and wastes in supply chains: A systematic review. *Waste Manag.* **2016**, *58*, 359–368. [CrossRef] [PubMed]
83. Pinto, A.T.; Schmidt, V.; Raimundo, S.A.; Raihmer, F. Use of ozone to control fungi in a cheese ripening room. *Acta Sci. Vet.* **2007**, *35*, 333–337. [CrossRef]
84. Serra, R.; Abrunhosa, L.; Kozakiewicz, Z.; Venâncio, A.; Lima, N. Use of ozone to reduce molds in a cheese ripening room. *J. Food Prot.* **2003**, *66*, 2355–2358. [CrossRef] [PubMed]
85. Varga, L.; Szigeti, J. Use of ozone in the dairy industry: A review. *Int. J. Dairy Technol.* **2016**, *69*, 157–168. [CrossRef]
86. Korukluoglu, M.; Sahan, Y.; Yigit, A. The fungicidal efficacy of various commercial disinfectants used in the food industry. *Ann. Microbiol.* **2006**, *56*, 325–330. [CrossRef]
87. Bundgaard-Nielsen, K.; Nielsen, P.V. Fungicidal effect of 15 disinfectants against 25 fungal contaminants commonly found in bread and cheese manufacturing. *J. Food Prot.* **1995**, *59*, 268–275. [CrossRef]
88. Martin, H.; Maris, P. Synergism between hydrogen peroxide and seventeen acids against five agri-food-borne fungi and one yeast strain. *J. Appl. Microbiol.* **2012**, *113*, 1451–1460. [CrossRef] [PubMed]
89. Lado, B.H.; Yousef, A.E. Alternative food-preservation technologies: Efficacy and mechanisms. *Microbes Infect.* **2002**, *4*, 433–440. [CrossRef]
90. European Union Council Directive 92/46/EEC of 16 June 1992 Laying Down the Health Rules for the Production and Placing on the Market of Raw Milk, Heat-Treated Milk and Milk-Based Products. Available online: <https://www.fsvps.ru/fsvps-docs/ru/usefulinf/files/es9246.pdf> (accessed on 27 July 2017).
91. Sakkas, L.; Moutafi, A.; Moschopoulou, E.; Moatsou, G. Assessment of treatment of various types of milk. *Food Chem.* **2014**, *15*, 293–301. [CrossRef] [PubMed]
92. Houbraken, J.; Samson, R.A.; Frisvad, J.C. Byssosclamyces: Significance of heat resistance and mycotoxin production. In *Advances in Food Mycology*; Hocking, A.D., Pitt, J.I., Samson, R.A., Thrane, U., Eds.; Springer: New York, NY, USA, 2005; pp. 211–224.
93. Scaramuzza, N.; Berni, E. Heat-resistance of *Hamigera avellanea* and *Thermoascus crustaceus* isolated from pasteurized acid products. *Int. J. Food Microbiol.* **2014**, *169*, 63–68. [CrossRef] [PubMed]
94. Salomão, B.C.M.; Slongo, A.P.; Aragão, G.M.F. Heat resistance of *Neosartorya fischeri* in various juices. *LWT Food Sci. Technol.* **2007**, *40*, 676–680. [CrossRef]
95. Trujillo, A.J.; Capellas, M.; Saldo, J.; Gervilla, R.; Guamis, B. Applications of high-hydrostatic pressure on milk and dairy products: A review. *Innov. Food Sci. Emerg. Technol.* **2002**, *3*, 295–307. [CrossRef]
96. Mainville, I.; Montpetit, D.; Durand, N.; Farnworth, E.R. Deactivating the bacteria and yeast in kefir using heat treatment, irradiation and high pressure. *Int. Dairy J.* **2001**, *11*, 45–49. [CrossRef]



97. Reps, A.; Krzyewska, A.; Laniewska-Moroz, L.; Iwanczak, M. Effect of high pressure on microflora of kefir. In Proceedings of the First International Conference on High Pressure Bioscience and Biotechnology HPBB-2000, Kyoto, Japan, 26–30 November 2000; Volume 19, pp. 1–652. [[CrossRef](#)]
98. Evert-Arriagada, K.; Hernández-Herrero, M.M.; Guamis, B.; Trujillo, A.J. Commercial application of high-pressure processing for increasing starter-free fresh cheese shelf-life. *Food Sci. Technol.* **2014**, *55*, 498–505. [[CrossRef](#)]
99. Ravishankar, S.; Zhang, H.; Kempkes, M.L. Pulsed electric fields. *Food Sci. Technol. Int.* **2008**, *14*, 429–432. [[CrossRef](#)]
100. Buckow, R.; Scott Chandry, P.; Ng, S.Y.; McAuley, C.M.; Swanson, B.G. Opportunities and challenges in pulse electric field processing of dairy products. *Int. Dairy J.* **2014**, *34*, 199–212. [[CrossRef](#)]
101. Wan, J.; Coventry, J.; Swiergon, P.; Sanguansri, P.; Viersteeg, C. Advances in innovative processing technologies for microbial inactivation and enhancement of food safety—Pulsed electric field and low temperature plasma. *Trends Food Sci. Technol.* **2009**, *20*, 414–424. [[CrossRef](#)]
102. Gougouli, M.; Kalantzi, K.; Beletsiotis, E.; Koutsoumanis, K.P. Development and application of predictive models for fungal growth as tools to improve quality control in yoghurt production. *Food Microbiol.* **2011**, *28*, 1453–1462. [[CrossRef](#)] [[PubMed](#)]
103. Sandhya. Modified atmosphere packaging of fresh produce: Current status and future needs. *Food Sci. Technol.* **2010**, *43*, 381–392. [[CrossRef](#)]
104. Ooraikul, B.; Stiles, M.E. *Modified Atmosphere Packaging of Food*; Ellis Horwood Publishing: New York, NY, USA, 1991; ISBN 978-0-7476-0064-0.
105. Nguyen Van Long, N.; Joly, C.; Dantigny, P. Active packaging with antifungal activities. *Int. J. Food Microbiol.* **2016**, *220*, 73–90. [[CrossRef](#)] [[PubMed](#)]
106. Rodriguez-Aguilera, R.; Oliveira, J.C.; Montanez, J.C.; Mahajan, P.V. Effect of modified atmosphere packaging on quality factors and shelf-life of surface mould ripened cheese: Part II varying storage temperature. *LWT Food Sci. Technol.* **2011**, *44*, 330–336. [[CrossRef](#)]
107. Sing, P.; Wani, A.A.; Karim, A.A.; Langowski, H.-C. The use of carbon dioxide in the processing and packaging of milk and dairy products: A review. *Int. J. Dairy Technol.* **2012**, *65*, 161–177. [[CrossRef](#)]
108. Hotchkiss, J.H.; Werner, B.G.; Lee, E.Y.C. Addition of carbon dioxide to dairy products to improve quality: A comprehensive review. *Compr. Rev. Food Sci. Food Saf.* **2006**, *5*, 158–168. [[CrossRef](#)]
109. Gonzalez-Fandos, E.; Sanz, S.; Olarte, C. Microbiological, physicochemical and sensory characteristics of Cameros cheese packaged under modified atmospheres. *Food Microbiol.* **2000**, *17*, 407–414. [[CrossRef](#)]
110. Taniwaki, M.H.; Hocking, A.D.; Pitt, J.I.; Fleet, G.H. Growth and mycotoxin production by fungi in atmosphere containing 80% carbon dioxide and 20% oxygen. *Int. J. Food Microbiol.* **2010**, *143*, 218–225. [[CrossRef](#)] [[PubMed](#)]
111. Khoshgozaran, S.; Azizi, M.H.; Bagheripoor-Fallah, N. Evaluating the effect of modified atmosphere packaging on cheese characteristics: A review. *Dairy Sci. Technol.* **2012**, *92*, 1–24. [[CrossRef](#)]
112. Taniwaki, M.H.; Hocking, A.D.; Pitt, J.I.; Fleet, G.H. Growth and mycotoxin production by food spoilage under high carbon dioxide and low oxygen atmospheres. *Int. J. Food Microbiol.* **2009**, *132*, 100–108. [[CrossRef](#)] [[PubMed](#)]
113. Dermiki, M.; Ntzimani, A.; Badeka, A.; Savvaidis, I.N.; Kontominas, M.G. Shelf-life extension and quality attributes of the whey cheese “Myzithra Kalathaki” using modified atmosphere packaging. *LWT Food Sci. Technol.* **2008**, *41*, 284–294. [[CrossRef](#)]
114. Eliot, S.C.; Vuilleumard, J.-C.; Edmond, J.-P. Stability of shredded Mozzarella cheese under modified atmospheres. *J. Food Sci.* **1998**, *63*, 1075–1080. [[CrossRef](#)]
115. Fedio, W.M.; Macleod, A.; Ozimek, L. The effect of modified atmosphere packaging on the growth of microorganisms in cottage cheese. *Milchwissenschaft* **1994**, *49*, 622–629.
116. Genigeorgis, C. Microbial and safety implications of the use of modified atmospheres to extend the storage life of fresh meat and fish. *Int. J. Food Microbiol.* **1985**, *1*, 237–251. [[CrossRef](#)]
117. Phillips, C.A. Review: Modified atmosphere packaging and its effects on the microbiological quality and safety of produce. *Int. J. Food Sci. Technol.* **1996**, *31*, 463–479. [[CrossRef](#)]
118. Rodriguez-Aguilera, R.; Oliveira, J.C.; Montanez, J.C.; Mahajan, P.V. Effect of modified atmosphere packaging on quality factors and shelf-life of surface mould ripened cheese: Part I constant temperature. *LWT Food Sci. Technol.* **2011**, *44*, 337–342. [[CrossRef](#)]

119. Blocher, J.C.; Busta, F.F. Multiple modes of inhibition of spore germination and outgrowth by reduced pH and sorbate. *J. Appl. Bacteriol.* **1985**, *59*, 469–478. [[CrossRef](#)] [[PubMed](#)]
120. Guynot, M.E.; Ramos, A.J.; Sanchis, V.; Marín, S. Study of benzoate, propionate, and sorbate salts as mould spoilage inhibitors on intermediate moisture bakery products of low pH (4.5–5.5). *Int. J. Food Microbiol.* **2005**, *101*, 161–168. [[CrossRef](#)] [[PubMed](#)]
121. Huang, Y.; Wilson, M.; Chapman, B.; Hocking, A.D. Evaluation of the efficacy of four weak acids as antifungal preservatives in low-acid intermediate moisture model food systems. *Food Microbiol.* **2009**, *27*, 6–33. [[CrossRef](#)] [[PubMed](#)]
122. Stratford, M.; Nebe-von-Caron, G.; Steels, H.; Novodvorska, M.; Ueckert, J.; Archer, D.B. Weak-acid preservatives: pH and proton movements in the yeast *Saccharomyces cerevisiae*. *Int. J. Food Microbiol.* **2013**, *161*, 164–171. [[CrossRef](#)] [[PubMed](#)]
123. Fajardo, P.; Martins, J.T.; Fuciños, C.; Pastrana, L.; Teixeira, J.A.; Vicente, A.A. Evaluation of a chitosan-based edible film as carrier of natamycin to improve the storability of Saloio cheese. *J. Food Eng.* **2010**, *101*, 349–356. [[CrossRef](#)]
124. EFSA. Scientific Opinion on the use of Natamycin (E 235) as a food additive. EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS). *EFSA J.* **2009**, *7*, 1412. [[CrossRef](#)]
125. FDA (2015). ICFSAN/Office of Food Additive Safety. GRAS Notice Natamycin. Published: 6 November 2015. Response Letter GRAS Notice No. GRN 000578. Available online: <https://www.fda.gov/Food/IngredientsPackagingLabeling/GRAS/NoticeInventory/ucm484505.htm> (accessed on 27 July 2017).
126. Brandão, L.R.; Rosa, J.C.C.; Nicoli, J.R.; Almeida, M.V.S.; do Carmo, A.P.; Queiros, H.T.; Castro, I.M. Investigating acid stress response in different *Saccharomyces* strains. *J. Mycol.* **2014**, *12*, 1–9. [[CrossRef](#)]
127. Ullah, A.; Chandrasekaran, G.; Brul, S.; Smits, J.G. Yeast adaptation to weak acids prevents futile energy expenditure. *Front. Microbiol.* **2013**, *4*, 1–10. [[CrossRef](#)] [[PubMed](#)]
128. Casas, E.; Ancos, B.; Valderrama, M.J.; Cano, P.; Peinado, J.M. Pentadiene production from potassium sorbate by osmotolerant yeasts. *Int. J. Food Microbiol.* **2004**, *94*, 93–96. [[CrossRef](#)] [[PubMed](#)]
129. Plumridge, A.; Stratford, M.; Lowe, K.C.; Archer, D.B. The weak-acid preservative sorbic acid is decarboxylated and detoxified by a phynylacrylic acid decarboxylase, PadA1, in the spoilage mold *Aspergillus niger*. *Appl. Environ. Microbiol.* **2008**, *74*, 550–552. [[CrossRef](#)] [[PubMed](#)]
130. Kallinteri, L.D.; Kostoula, O.K.; Savvaidis, I.N. Efficacy of nisin and/or natamycin to improve the shelf-life of Galotyri cheese. *Food Microbiol.* **2013**, *36*, 176–181. [[CrossRef](#)] [[PubMed](#)]
131. De Oliveira, T.M.; De Fátima Ferreira Soares, N.; Pereira, R.M.; De Freitas Fraga, K. Development and evaluation of antimicrobial natamycin-incorporated film in Gorgonzola cheese conservation. *Packag. Technol. Sci.* **2007**, *20*, 147–153. [[CrossRef](#)]
132. Ollé Resa, C.P.; Gerschenson, L.N.; Jagus, R.J. Natamycin and nisin supported on starch edible films for controlling mixed growth on model systems and Port Salut cheese. *Food Control* **2014**, *44*, 146–151. [[CrossRef](#)]
133. Ollé Resa, C.P.; Jagus, R.J.; Gerschenson, L.N. Natamycin efficiency for controlling yeast growth in models systems and on cheese surfaces. *Food Control* **2014**, *35*, 101–108. [[CrossRef](#)]
134. WHO. *Safety Evaluation of Certain Food Additives and Contaminants; Natamycin (Pimaricin)*; WHO Food Additives Series 48; WHO: Geneva, Switzerland, 2002. Available online: [http://apps.who.int/iris/bitstream/10665/42578/1/WHO\\_TRS\\_909.pdf](http://apps.who.int/iris/bitstream/10665/42578/1/WHO_TRS_909.pdf) (accessed on 27 July 2017).
135. Van Leeuwen, M.R.; De Boer, W.; Smant, W.; Dijksterhuis, J. Filipin is a reliable in situ marker of ergosterol in the plasma membrane of germinating conidia (spores) of *Penicillium discolor* and stains intensively at the site of germ tube formation. *J. Microbiol. Methods* **2008**, *74*, 64–73. [[CrossRef](#)] [[PubMed](#)]
136. Van Leeuwen, M.R.; Van Doorn, T.; Golovina, E.A.; Stark, J.; Dijksterhuis, J. Water and air-distributed conidia exhibit differences in sterol content and cytoplasmic microviscosity. *Appl. Environ. Microbiol.* **2010**, *67*, 366–369. [[CrossRef](#)] [[PubMed](#)]
137. Stern, G.A. In vitro antibiotic synergism against ocular fungal isolates. *Am. J. Ophthalmol.* **1978**, *86*, 359–367. [[CrossRef](#)]
138. Streekstra, H.; Verkennis, A.E.E.; Jacobs, R.; Dekker, A.; Stark, J.; Dijksterhuis, J. Fungal strain and the development of tolerance against natamycin. *Int. J. Food Microbiol.* **2016**, *238*, 15–22. [[CrossRef](#)] [[PubMed](#)]

139. Joint FAO/WHO Expert Committee on Food Additives. *Evaluation of Certain Food Additives and Contaminants: 67th Report of the Joint FAO/WHO Expert Committee on Food Additives*; WHO Technical Report Series; WHO: Geneva, Switzerland, 2007; p. 25. Available online: <http://www.who.int/ipcs/publications/jecfa/reports/trs940.pdf> (accessed on 27 July 2017).
140. Bourdichon, F.; Casaregola, S.; Farrokhi, C.; Frisvad, J.C.; Gerds, M.L.; Hammes, W.P.; Harnett, J.; Huys, G.; Laulund, S.; Ouwehand, A.; et al. Food fermentations: Microorganisms with technological beneficial use. *Int. J. Food Microbiol.* **2012**, *154*, 87–97. [[CrossRef](#)] [[PubMed](#)]
141. Caplice, E.; Fitzgerald, G.F. Food fermentations: Role of microorganisms in food production and preservation. *Int. J. Food Microbiol.* **1999**, *50*, 131–149. [[CrossRef](#)]
142. Brimelow, C.J.B.; Collins-Thompson, D. Food preservatives. *Food Control* **1992**, *3*, 218–219. [[CrossRef](#)]
143. Elsser-Gravesen, D.; Elsser-Gravesen, A. Biopreservatives. In *Biotechnology of Food and Feed Additives*; Zorn, H., Cermak, P., Eds.; Springer: Berlin, Germany, 2014; ISBN 978-3-662-43761-2.
144. Stiles, M.E. Biopreservation by lactic acid bacteria. *Antonie van Leeuwenhoek* **1996**, *70*, 331–345. [[CrossRef](#)] [[PubMed](#)]
145. Holzapfel, W.H.; Geisen, R.; Schillinger, U. Biological preservation of foods with reference to protective cultures, bacteriocins and food-grade enzymes. *Int. J. Food Microbiol.* **1995**, *24*, 343–362. [[CrossRef](#)]
146. Magnusson, J.; Schnurer, J. *Lactobacillus coryniformis* subs. *coryniformis* strain Si3 produces a broad-spectrum proteinaceous antifungal compound. *Appl. Environ. Microbiol.* **2001**, *67*, 1–5. [[CrossRef](#)] [[PubMed](#)]
147. Lassois, L.; de Lapeyre de Bellaire, L.; Jijakli, M.H. Biological control of crown rot of bananas with *Pichia anomala* strain K and *Candida oleophila* strain O. *Biol. Control* **2008**, *45*, 410–418. [[CrossRef](#)]
148. Zhang, H.; Wang, L.; Ma, L.; Dong, Y.; Jiang, S.; Xu, B.; Zheng, X. Biocontrol of major postharvest pathogens on apple using *Rhodotorula glutinis* and its effects on postharvest quality parameters. *Biol. Control* **2009**, *48*, 79–83. [[CrossRef](#)]
149. Carr, F.J.; Chill, D.; Maida, N. The lactic acid bacteria: A literature survey. *Crit. Rev. Microbiol.* **2002**, *28*, 281–370. [[CrossRef](#)] [[PubMed](#)]
150. Schwenninger, S.M.; Meile, L. A mixed culture of *Propionibacterium jensenii* and *Lactobacillus paracasei* subs. *paracasei* inhibits food spoilage yeasts. *Syst. Appl. Microbiol.* **2004**, *27*, 229–237. [[CrossRef](#)] [[PubMed](#)]
151. Aljewicz, M.; Cichosz, G. Protective effects of *Lactobacillus* cultures in Dutch-type cheese like products. *LWT Food Sci. Technol.* **2015**, *63*, 52–56. [[CrossRef](#)]
152. Aunbjerg, S.D.; Honoré, A.H.; Marcussen, J.; Ebrahimi, P.; Vogensen, F.K.; Benfeldt, C.; Skov, T.; Knøchel, S. Contribution of volatiles to the antifungal effect of *Lactobacillus paracasei* in defined medium and yoghurt. *Int. J. Food Microbiol.* **2015**, *194*, 46–53. [[CrossRef](#)] [[PubMed](#)]
153. Crowley, S.; Mahony, J.; van Sinderen, D. Current perspectives on antifungal lactic acid bacteria as natural bio-preservatives. *Trends Food Sci. Technol.* **2013**, *33*, 93–109. [[CrossRef](#)]
154. Inglin, R.C.; Stevens, M.J.A.; Meile, L.; Lacroix, C.; Meile, L. High-throughput screening assays for antibacterial and antifungal activities of *Lactobacillus* species. *J. Microbiol. Methods* **2015**, *114*, 26–29. [[CrossRef](#)] [[PubMed](#)]
155. Stiles, J.; Penkar, S.; Plocková, M.; Chumchalová, J.; Bullerman, L.B. Antifungal activity of sodium acetate and *Lactobacillus rhamnosus*. *J. Food Prot.* **2002**, *65*, 1188–1191. [[CrossRef](#)] [[PubMed](#)]
156. Le Lay, C.; Mounier, J.; Vasseur, V.; Weill, A.; Le Blay, G.; Barbier, G.; Coton, E. In vitro and in situ screening of lactic acid bacteria and propionibacteria antifungal activities against bakery product spoilage molds. *Food Control* **2016**, *60*, 247–255. [[CrossRef](#)]
157. Delavenne, E.; Mounier, J.; Deniel, F.; Barbier, G.; Le Blay, G. Biodiversity of antifungal lactic acid bacteria isolated from raw milk samples from cow, ewe and goat over one-year period. *Int. J. Food. Microbiol.* **2012**, *155*, 185–190. [[CrossRef](#)] [[PubMed](#)]
158. Fernandez, B.; Vimon, A.; Desfossés-Foucault, E.; Daga, M.; Arora, G.; Fliss, I. Antifungal activity of lactic and propionic acid bacteria and their potential as protective culture in cottage cheese. *Food Control* **2017**, *78*, 350–356. [[CrossRef](#)]
159. Axelsson, L.; Chung, T.C.; Debrogosz, W.J.; Lindgren, S. Production of a broad spectrum antimicrobial substance by *Lactobacillus reuteri*. *Microb. Ecol. Health Dis.* **1989**, *2*, 131–136. [[CrossRef](#)]
160. Dagnas, S.; Gauvry, E.; Onno, B.; Membré, J.M. Quantifying effect of lactic, acetic, and propionic acids on growth of molds isolated from spoiled bakery products. *J. Food Prot.* **2015**, *78*, 1689–1698. [[CrossRef](#)] [[PubMed](#)]

161. Coallier-Ascah, J.; Idziak, E.S. Interaction between *Streptococcus lactis* and *Aspergillus flavus* on production of aflatoxin. *Appl. Environ. Microbiol.* **1985**, *49*, 163–167. [[CrossRef](#)] [[PubMed](#)]
162. Luchese, R.H.; Harrigan, W.F. Growth of, and aflatoxin production by *Aspergillus paraciticus* when in the presence of either *Lactococcus lactis* or lactic acid and at different initial pH values. *J. Bacteriol.* **1990**, *69*, 512–519. [[CrossRef](#)]
163. Batish, V.K.; Roy, U.; Lal, R.; Grower, S. Antifungal attributes of lactic acid bacteria—A review. *Crit. Rev. Biotechnol.* **1997**, *17*, 209–225. [[CrossRef](#)] [[PubMed](#)]
164. Roy, U.; Batish, V.K.; Grover, S.; Neelakantan, S. Production of antifungal substance by *Lactococcus lactis* subsp. *lactis* CHD-28.3. *Int. J. Food Microbiol.* **1996**, *32*, 27–34. [[CrossRef](#)]
165. Florianowicz, T. Antifungal activity of some microorganisms against *Penicillium expansum*. *Eur. Food Res. Technol.* **2001**, *212*, 282–286. [[CrossRef](#)]
166. Voulgari, K.; Hatzikamari, M.; Delepoglou, A.; Georgakipoulos, P.; Litopoulou-Tzanetaki, E.; Tzanetakis, N. Antifungal activity of non-starter lactic acid bacteria isolates from dairy products. *Food Control* **2010**, *21*, 136–142. [[CrossRef](#)]
167. Garofalo, C.; Zannini, E.; Aquilanti, L.; Silvestri, G.; Fierro, O.; Picariello, G.; Clementi, F. Selection of sourdough lactobacilli with antifungal activity for use as biopreservatives in bakery products. *J. Agric. Food Chem.* **2012**, *60*, 7719–7728. [[CrossRef](#)] [[PubMed](#)]
168. Coda, R.; Rizello, C.G.; Nigri, F.; De Angelis, M.; Arnault, P.; Gobbetti, M. Long-term fungi inhibitory activity of water-soluble extract from *Phaseolus vulgaris* cv. pinto and sourdough lactic acid bacteria during bread storage. *Appl. Environ. Microbiol.* **2008**, *74*, 7391–7398. [[CrossRef](#)] [[PubMed](#)]
169. Gerez, C.L.; Torres, M.J.; Font de Valdez, G.; Rollán, G. Control of spoilage fungi by lactic acid bacteria. *Biol. Control* **2013**, *64*, 231–237. [[CrossRef](#)]
170. Mauch, A.; Dal Bello, F.; Coffey, A.; Arendt, E.K. The use of *Lactobacillus brevis* PS1 to in vitro inhibit the outgrowth of *Fusarium culmorum* and other common *Fusarium* species found on barley. *Int. J. Food Microbiol.* **2010**, *141*, 116–121. [[CrossRef](#)] [[PubMed](#)]
171. Falguni, P.; Shilpa, V.; Mann, B. Production of proteinaceous antifungal substances from *Lactobacillus brevis* NCDC. *Int. J. Dairy Technol.* **2010**, *63*, 70–76. [[CrossRef](#)]
172. Gourama, H. Inhibition of growth and mycotoxin production of *Penicillium* by *Lactobacillus* species. *LWT Food Sci. Technol.* **1997**, *30*, 279–283. [[CrossRef](#)]
173. Li, H.; Liu, L.; Zhang, S.; Uluko, H.; Cui, W.; Lv, J. Potential use of *Lactobacillus casei* AST18 as bioprotective culture in yoghurt. *Food Control* **2013**, *34*, 675–680. [[CrossRef](#)]
174. Atanassova, M.R.; Fernández-Otero, C.; Rodríguez-Alonso, P.; Fernández-No, I.C.; Garabal, J.I.; Centeno, J.A. Characterization of yeasts isolated from artisanal short-ripened cows' cheeses produced in Galicia (NW Spain). *Food Microbiol.* **2016**, *53*, 172–181. [[CrossRef](#)] [[PubMed](#)]
175. Ryan, L.A.; Zannini, E.; Dal Bello, F.; Pawlowska, A.; Koehler, P.; Arendt, E.K. *Lactobacillus amylovorus* DSM 19280 as a novel food-grade antifungal agent for bakery products. *Int. J. Food Microbiol.* **2011**, *146*, 276–283. [[CrossRef](#)]
176. Lynch, K.M.; Pawlowska, A.M.; Brosnan, B.; Coffey, A.; Zannini, E.; Furey, A.; McSweeney, P.L.H.; Waters, D.M.; Arendt, E.K. Application of *Lactobacillus amylovorus* as an antifungal adjunct to extend the shelf-life of Cheddar cheese. *Int. J. Food Microbiol.* **2014**, *34*, 167–173. [[CrossRef](#)]
177. Corsetti, A.; Gobetti, M.; Rossi, J.; Damiani, P. Antimould activity of sourdough lactic acid bacteria: Identification of a mixture of organic acids produced by *Lactobacillus sanfrancisco* CB1. *Appl. Microbiol. Biotechnol.* **1998**, *50*, 253–256. [[CrossRef](#)] [[PubMed](#)]
178. Corsetti, A.; Settani, L.; van Sinderen, D.; Felis, G.; Dellaglio, F.; Gobetti, M. *Lactobacillus rossii* sp. nov., isolated from wheat sourdough. *Int. J. Syst. Evol. Microbiol.* **2004**, *55*, 35–40. [[CrossRef](#)]
179. Muhialdin, B.J.; Hassan, Z.; Sadon, S.K.; Zulkifli, N.A.; Azfar, A.A. Effect of pH and heat treatment on antifungal activity of *Lactobacillus fermentum* TE007, *Lactobacillus pentosus* G004 and *Pediococcus pentosaceus* TE010. *Innov. Romanian Food Biotechnol.* **2011**, *8*, 41–53. [[CrossRef](#)]
180. Coloretto, F.; Carri, S.; Armaforte, E.; Chiavari, C.; Grazia, L.; Zambonelli, C. Antifungal activity of lactobacilli isolated from salami. *FEMS Microbiol. Lett.* **2007**, *271*, 245–250. [[CrossRef](#)] [[PubMed](#)]
181. Guo, J.; Mauch, A.; Galle, S.; Murphy, P.; Arendt, E.K.; Coffey, A. Inhibition of growth of *Trichophyton tonsurans* by *Lactobacillus reuteri*. *J. Appl. Microbiol.* **2011**, *111*, 474–483. [[CrossRef](#)] [[PubMed](#)]



182. Niku-Paavola, M.L.; Laitila, A.; Mattila-Sandholm, T.; Haikara, A. New types of antimicrobial compounds produced by *Lactobacillus plantarum*. *J. Appl. Microbiol.* **1999**, *86*, 29–35. [[CrossRef](#)] [[PubMed](#)]
183. Dal Bello, F.; Clarke, C.; Ryan, L.; Ulmer, H.; Schober, T.; Strom, K.; Sjögren, J.; van Sinderen, D.; Schnürer, J.; Arendt, E.K. Improvement of the quality and shelf life of wheat bread by fermentation with the antifungal strain *Lactobacillus plantarum* FST 1.7. *J. Cereal Sci.* **2007**, *45*, 309–318. [[CrossRef](#)]
184. Lavermicocca, P.; Valerio, F.; Visconti, A. Antifungal activity of phenyllactic acid against molds isolated from Bakery products. *Appl. Environ. Microbiol.* **2003**, *69*, 634–640. [[CrossRef](#)] [[PubMed](#)]
185. Ström, K.; Sjögren, J.; Broberg, A.; Schnürer, J. *Lactobacillus plantarum* MiLAB 393 produces the antifungal cyclic dipeptides cyclo (L-Phe-L-Pro) and cyclo (L-Phe-trans-4-OH-L-Pro) and 3-phenyllactic acid. *Appl. Environ. Microbiol.* **2002**, *68*, 4322–4327. [[CrossRef](#)] [[PubMed](#)]
186. Sjögren, J.; Magnusson, J.; Broberg, A.; Schnürer, J.; Kenne, L. Antifungal 3- hydroxy fatty acids from *Lactobacillus plantarum* MiLAB 14. *Appl. Environ. Microbiol.* **2003**, *69*, 7554–7557. [[CrossRef](#)] [[PubMed](#)]
187. Sathe, S.J.; Nawani, N.N.; Dhakephalkar, P.K.; Kapadnis, B.P. Antifungal lactic acid bacteria with potential to prolong shelf-life of fresh vegetables. *J. Appl. Microbiol.* **2007**, *103*, 2622–2628. [[CrossRef](#)] [[PubMed](#)]
188. Yang, E.J.; Chang, H.C. Purification of a new antifungal compound produced by *Lactobacillus plantarum* AF1 isolated from kimchi. *Int. J. Food Microbiol.* **2010**, *139*, 56–63. [[CrossRef](#)] [[PubMed](#)]
189. Prema, P.; Smila, D.; Palavesan, A.; Immanuel, G. Production and characterization of an antifungal compound (3-phenyllactic acid) produced by *Lactobacillus plantarum* strain. *Food Bioprocess Technol.* **2008**, *3*, 379–386. [[CrossRef](#)]
190. Rouse, S.; Harnett, D.; Vaughan, A.; Sinderen, D.V. Lactic acid bacteria with potential to eliminate fungal spoilage in foods. *J. Appl. Microbiol.* **2008**, *104*, 915–923. [[CrossRef](#)] [[PubMed](#)]
191. Ndagano, D.; Lamoureux, T.; Dortu, C.; Vandermoten, S.; Thonart, P. Antifungal activity of 2 lactic acid bacteria of the *Weissella* Genus Isolated from Food. *J. Food Sci.* **2011**, *76*, 305–311. [[CrossRef](#)] [[PubMed](#)]
192. Wang, H.; Sun, Y.; Chen, C.; Sun, Z.; Zhou, Y.; Shen, F. Genome shuffling of *Lactobacillus plantarum* for improving antifungal activity. *Food Control* **2013**, *32*, 341–347. [[CrossRef](#)]
193. Laitila, A.; Alakomi, H.L.; Raaska, L.; Mattila-Sandholm, T.; Haikara, A. Antifungal activities of two *Lactobacillus plantarum* strains against *Fusarium* moulds in vitro and in malting of barley. *J. Appl. Microbiol.* **2002**, *93*, 566–576. [[CrossRef](#)]
194. Digaitiene, A.; Hansen, A.; Juodeikiene, G.; Eidukonyte, D.; Josephsen, J. Lactic acid bacteria isolated from rye sourdoughs produce bacteriocin-like inhibitory substances active against *Bacillus subtilis* and fungi. *J. Appl. Microbiol.* **2012**, *112*, 732–742. [[CrossRef](#)] [[PubMed](#)]
195. Lind, H.; Jonsson, H.; Schnürer, J. Antifungal effect of dairy propionibacteria—contribution of organic acids. *Int. J. Food Microbiol.* **2003**, *98*, 157–165. [[CrossRef](#)] [[PubMed](#)]
196. Mieszkina, S.; Hymery, N.; Debaets, S.; Coton, E.; Le Blay, G.; Valence, F.; Mounier, J. Action mechanism involved in the bioprotective effect of *Lactobacillus harbinensis* K.V9.3.1Np against *Yarrowia lipolytica* in fermented milk. *Int. J. Food Microbiol.* **2017**, *248*, 47–55. [[CrossRef](#)] [[PubMed](#)]
197. Al-Zoreky, N.; Ayres, J.W.; Sandine, W.E. Antimicrobial activity of Microgard against food spoilage and pathogenic microorganisms. *J. Dairy Sci.* **1991**, *74*, 758–763. [[CrossRef](#)]
198. Daeschel, A.M. Antimicrobial substances from lactic acid bacteria for use as food preservatives. *Food Technol.* **1989**, *43*, 91–94. [[CrossRef](#)]

